

10/031146
JC10 Rec'd PCT/PTO 17 JAN 2002

TRANSMITTAL LETTER TO THE UNITED STATES

ATTORNEY'S DOCKET NUMBER 50915

DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

INTERNATIONAL APPLICATION NO.	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED
PCT/EP00/07253	27 July 2000	27 July 1999 18 November 1999 22 March 2000
TITLE OF INVENTION: NOVEL CYTOCHROME P450 MONOOXYGENASES AND THEIR USE FOR OXIDIZING ORGANIC COMPOUNDS		
APPLICANT(S) FOR DO/EO/US Bernhard HAUER, Juergen PLEISS, Ulrich SCHWANEBERG, Jutta SCHMITT, Markus FISCHER Rolf SCHMID, Qing-shan LI, Sabine Lutz-WAHL, Daniel APPEL		

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. /X/ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
 2. / / This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
 3. /X/ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
 4. /x / A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
 5. /X/ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. /X/ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. / / has been transmitted by the International Bureau.
 - c. / / is not required, as the application was filed in the United States Receiving Office (RO/USO).
 6. /X/ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
 7. /X / Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
 - a. / X / are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. / / have been transmitted by the International Bureau.
 - c. / / have not been made; however, the time limit for making such amendments has NOT expired.
 - d. / / have not been made and will not be made.
 8. / X / A translation of the amendments to the claims under PCT Article 19(35 U.S.C. 371(c)(3)).
 9. / X / An oath or declaration of the inventor(s)(35 U.S.C. 171(c)(4)).
 10. / / A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).
- Items 11. to 16. below concern other document(s) or information included:
11. / / An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
 12. / X / An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
 13. /X / A FIRST preliminary amendment.
/ / A SECOND or SUBSEQUENT preliminary amendment.
 14. / / A substitute specification.
 15. / / A change of power of attorney and/or address letter.
 16. /x / Other items or information.
International Search Report
International Preliminary Examination Report

U.S. Appln. No. (If Known) INTERNATIONAL APPLN. NO.
PCT/EP00/ 07253


ATTORNEY'S DOCKET NO.
50915

		CALCULATIONS	PTO USE ONLY
17. /X/ The following fees are submitted			
BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):			
Search Report has been prepared by the			
EPO or JPO.....	\$890.00	890.00	
International preliminary examination fee paid to USPTO			
(37 CFR 1.482).....	\$710.00		
No international preliminary examination fee paid to			
USPTO (37 CFR 1.482) but international search fee paid			
to USPTO (37 CFR 1.445(a)(2)).....	\$740.00		
Neither international preliminary examination fee			
(37 CFR 1.482) nor international search fee			
(37 CFR 1.445(a)(2)) paid to USPTO	\$ 1,040.00		
International preliminary examination fee paid to			
USPTO (37 CFR 1.482) and all claims satisfied pro			
-visions of PCT Article 33(2)-(4).....	\$100.00		
ENTER APPROPRIATE BASIC FEE AMOUNT = \$		890.00	
Surcharge of \$130.00 for furnishing the oath or declaration			
later than // 20 // 30 months from the earliest			
claimed priority date (37 CFR 1.492(e)).			
Claims	Number Filed	Number Extra	Rate
Total Claims	23 -20	3	X\$18. 54.
Indep. Claims	2 -3		X\$84.
Multiple dependent claim(s)(if applicable)			+280.
TOTAL OF ABOVE CALCULATION		=	944.
Reduction of 1/2 for filing by small entity, if applicable.			
Verified Small Entity statement must also be filed			
(Note 37 CFR 1.9, 1.27, 1.28).			
SUBTOTAL		=	944.
Processing fee of \$130. for furnishing the English			
translation later than // 20 // 30 months from the			
earliest claimed priority date (37 CFR 1.492(f)). +			
TOTAL NATIONAL FEE		=	944 .
Fee for recording the enclosed assignment (37 CFR 1.21(h)).			
The assignment must be accompanied by an appropriate cover			
sheet (37 CFR 3.28, 3.31) \$40.00 per property =			
TOTAL FEES ENCLOSED		=	\$ 984.00
		Amount to be	
		refunded:	\$
		Charged	\$

- a./X/ A check in the amount of \$ 984.00 to cover the above fees is enclosed.
- b./ / Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed.
- c./X/ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 11-0345. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:
KEIL & WEINKAUF
1101 Connecticut Ave., N.W.
Washington, D. C. 20036


SIGNATURE
Herbert B. Keil
NAME
Registration No. 18,967

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of)
HAUER et al.) BOX PCT
)
International Application)
PCT/EP 00/07253)
)
Filed: July 27, 2000)
)

For: NOVEL CYTOCHROME P450 MONOOXYGENASES AND THEIR USE FOR THE
OXIDATION OF ORGANIC COMPOUNDS

PRELIMINARY AMENDMENT

Honorable Commissioner of
Patents and Trademarks
Washington, D.C. 20231

Sir:

Prior to examination, kindly amend the above-identified application as follows:

IN THE CLAIMS

Kindly amend the claims as shown on the attached sheets.

R E M A R K S


The claims were amended in the preliminary examination. The claims have been amended to eliminate multiple dependency and to place them in better form for U.S. filing. No new matter is included.

A clean copy of the claims is attached.

Favorable action is solicited.

Respectfully submitted,

KEIL & WEINKAUF


Herbert B. Keil
Reg. No. 18,967

1101 Connecticut Ave., N.W.
Washington, D.C. 20036

(202)659-0100

CLEAN VERSION OF AMENDED CLAIMS - 50915

4. A nucleic acid sequence coding for a monooxygenase according to claim 1.
11. A process as claimed in claim 9, where the monooxygenase is derived from cytochrome P450 monooxygenase BM-3 from *Bacillus megaterium* having an amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43. 48-52. 67-70, 330-335, 352-356, 73-82 and 86-88.
13. A process for microbiological oxidation of optionally substituted mono- or polynuclear aromatics, straight-chain or branched alkanes or alkenes, or optionally substituted cycloalkanes or cycloalkenes, which comprises
 - a1) culturing the recombinant cytochrome P450-producing microorganism as claimed in claim 7 in a culture medium, in the presence of an exogenous or intermediately formed substrate; or
 - a2) incubating a substrate-containing reaction medium with a cytochrome P450 monooxygenase derived from cytochrome P450 monooxygenase BM-3 from *Bacillus megaterium* having an amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43. 48-52. 67-70, 330-335, 352-356, 73-82 and 86-88; and
 - b) isolating the oxidation product formed or a secondary product thereof from the medium;where the monooxygenase mutant Phe87Val is not excluded.

CLEAN VERSION OF AMENDED CLAIMS - 50915

15. (cancel)
16. A process as claimed in claim 13, where the cytochrome P450 monooxygenase has at least one of the following mono- or polyamino acid substitutions:
 - a) Phe87Val;
 - b) Phe87Val, Leu188Gln; or
 - c) Phe87Val, Leu188Gln, Ala74Gly.
17. A process as claimed in claim 9, wherein, as exogenous substrate, at least one compound selected from the groups a) to d) of compounds defined above is added to a medium and the oxidation is carried out by enzymatic reaction of the substrate-containing medium in the presence of oxygen at a temperature of approximately 20 to 40°C and a pH of approximately 6 to 9, where the substrate-containing medium additionally contains an approximately 10- to 100-fold molar excess of reduction equivalents based on the substrate.
19. A process for the microbiological production of indigo and/or indixubin, which comprises
 - a1) culturing a recombinant microorganism which produces an indole-oxidizing cytochrome P450 in a culture medium, in the presence of exogenous or intermediately formed indole; or
 - a2) incubating an indole-containing reaction medium with an indole-oxidizing cytochrome P450 monooxygenase; and
 - b) isolating the oxidation product formed or a secondary product thereof

CLEAN VERSION OF AMENDED CLAIMS - 50915

from the medium.

22. A process as claimed in claim 20 , where the monooxygenase is derived from cytochrome P450 monooxygenase BM-3 from *Bacillus megaterium* having an amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43, 48-52, 67-70, 330-335, 352-356, 73-82 and 86-88, including the substitution Phe87Val.
23. A process as claimed in claim 22, where the monooxygenase has at least one of the following mono- or polyamino acid substitutions:
 - a) Phe87Val;
 - b) Phe87Val, Leu188Gln; or
 - c) Phe87Val, Leu188Gln, Ala74Gly.
24. A bioreactor comprising the cytochrom P450 monooxygenase as claimed in claim 1 or a recombinant microorganism transformed by a vector comprising an expression construct comprising a nucleic acid sequence coding for the cytochrom P450 monooxygenase of claim 1 in immobilized form.
25. (cancel)
26. (cancel)

MARKED VERSION OF AMENDED CLAIMS - 50915

4. (amended) A nucleic acid sequence coding for a monooxygenase according to [one of the preceding claims] claim 1.
11. (amended) A process as claimed in claim 9 [or 10], where the monooxygenase is [a mutant as claimed in any of claims 1 to 3, including the mutant Phe87Val] derived from cytochrome P450 monooxygenase BM-3 from Bacillus megaterium having an amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43, 48-52, 67-70, 330-335, 352-356, 73-82 and 86-88.
13. (amended) A process for microbiological oxidation of [a compound as defined in claim 1 b), c) or d)] optionally substituted mono- or polynuclear aromatics, straight-chain or branched alkanes or alkenes, or optionally substituted cycloalkanes or cycloalkenes, which comprises
 - a1) culturing [a] the recombinant cytochrome P450-producing microorganism as claimed in claim 7 [or 8] in a culture medium, in the presence of an exogenous or intermediately formed substrate; or
 - a2) incubating a substrate-containing reaction medium with a cytochrome P450 monooxygenase [as claimed in any of claims 1 to 3] derived from cytochrome P450 monooxygenase BM-3 from Bacillus megaterium having an amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43, 48-52, 67-70, 330-335, 352-356, 73-82 and 86-

MARKED: VERSION OF AMENDED CLAIMS - 50915

15. (cancel)
16. (amended) A process as claimed in claim [15] 13, where the [mutant] cytochrome P450 monooxygenase has at least one of the following mono- or polyamino acid substitutions:
 - a) Phe87Val;
 - b) Phe87Val, Leu188Gln; or
 - c) Phe87Val, Leu188Gln, Ala74Gly.
17. (amended) A process as claimed in [any of claims] claim 9 [to 16], wherein, as exogenous substrate, at least one compound selected from the groups a) to d) of compounds defined above is added to a medium and the oxidation is carried out by enzymatic reaction of the substrate-containing medium in the presence of oxygen at a temperature of approximately 20 to 40°C and a pH of approximately 6 to 9, where the substrate-containing medium additionally contains an approximately 10- to 100-fold molar excess of reduction equivalents based on the substrate.
19. (amended) A process for the microbiological production of indigo and/or indixubin, which comprises
 - a1) culturing a recombinant microorganism which produces an indole-oxidizing cytochrome P450 in a culture medium, in the presence of exogenous or intermediately formed indole; or
 - a2) incubating an indole-containing reaction medium with an indole-oxidizing

MARKED VERSION OF AMENDED CLAIMS - 50915

- a1) culturing a recombinant microorganism which produces an indole-oxidizing cytochrome P450 in a culture medium, in the presence of exogenous or intermediately formed indole; or
 - a2) incubating an indole-containing reaction medium with an indole-oxidizing cytochrome P450 monooxygenase; and
 - b) isolating the oxidation product formed or a secondary product thereof from the medium[;].
22. (amended) A process as claimed in claim 20 [or 21], where the monooxygenase is [a mutant as claimed in any of claims 1 to 3] derived from cytochrome P450 monooxygenase BM-3 from Bacillus megaterium having an amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43, 48-52, 67-70, 330-335, 352-356, 73-82 and 86-88, including the [mutant] substitution Phe87Val.
23. (amended) A process as claimed in claim 22, where the [mutant] monooxygenase has at least one of the following mono- or polyamino acid substitutions:
- a) Phe87Val;
 - b) Phe87Val, Leu188Gln; or
 - c) Phe87Val, Leu188Gln, Ala74Gly.
24. (amended) A bioreactor comprising [an enzyme] the cytochrom P450 monooxygenase as claimed in [one of claims] claim 1 [to 3] or a recombinant

MARKED VERSION OF AMENDED CLAIMS - 50915

microorganism [as claimed in one of claims 7 or 8] transformed by a vector comprising an expression construct comprising a nucleic acid sequence coding for the cytochrom P450 monooxygenase of claim 1 in immobilized form.

25. (cancel)

26. (cancel)

CLAIMS AS FILED - 50915

1. A cytochrome P450 monooxygenase which is capable of at least one of the following reactions:
 - a) oxidation of optionally substituted N-, O- or S-heterocyclic mono- or polynuclear aromatic compounds;
 - b) oxidation of optionally substituted mono- or polynuclear aromatics;
 - c) oxidation of straight-chain or branched alkanes and alkenes;
 - d) oxidation of optionally substituted cycloalkanes and cycloalkenes;where the monooxygenase is derived from cytochrome P450 monooxygenase BM-3 from *Bacillus megaterium* having an amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43, 48-52, 67-70, 330-335, 352-356, 73-82 and 86-88; except the single mutant Phe87Val.
2. A monooxygenase as claimed in claim 1, which has at least one functional mutation in at least one of the sequence regions 73-82, 86-88 and 172-224.
3. A monooxygenase as claimed in claim 1, which has at least one of the following mono- or polyamino acid substitutions:
 - a) Phe87Val, Leu188Gln; or
 - b) Phe87Val, Leu188Gln, Ala74Gly;and functional equivalents thereof which are capable of at least one of the above oxidation reactions.

CLAIMS AS FILED - 50915

4. A nucleic acid sequence coding for a monooxygenase according to claim 1.
5. An expression construct comprising, under the genetic control of regulatory nucleic acid sequences, a coding sequence which comprises a nucleic acid sequence according to claim 4.
6. A vector comprising at least one expression construct according to claim 5.
7. A recombinant microorganism transformed by at least one vector as claimed in claim 6.
8. A microorganism as claimed in claim 7, selected from bacteria of the genus *Escherichia*.
9. A process for the microbiological oxidation of an N- or S-heterocyclic mono- or polynuclear aromatic compound which comprises
 - a1) culturing a recombinant microorganism which expresses a cytochrome P450 monooxygenase of bacterial origin in a culture medium, in the presence of an exogenous or intermediately formed substrate; or
 - a2) incubating a substrate-containing reaction medium with a cytochrome P450 monooxygenase of bacterial origin; and
 - b) isolating the oxidation product formed or a secondary product thereof from the medium.
10. A process as claimed in claim 9, wherein the exogenous or intermediately formed substrate is selected from optionally substituted - or S-heterocyclic mono- or polynuclear aromatic compounds.

CLAIMS AS FILED - 50915

11. A process as claimed in claim 9, where the monooxygenase is derived from cytochrome P450 monooxygenase BM-3 from *Bacillus megaterium* having an amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43, 48-52, 67-70, 330-335, 352-356, 73-82 and 86-88.
12. A process as claimed in claim 11, where the mutant has at least one of the following mono- or polyamino acid substitutions:
 - a) Phe87Val;
 - b) Phe87Val, Leu188Gln; or
 - c) Phe87Val, Leu188Gln, Ala74Gly.
13. A process for microbiological oxidation of optionally substituted mono- or polynuclear aromatics, straight-chain or branched alkanes or alkenes, or optionally substituted cycloalkanes or cycloalkenes, which comprises
 - a1) culturing the recombinant cytochrome P450-producing microorganism as claimed in claim 7 in a culture medium, in the presence of an exogenous or intermediately formed substrate; or
 - a2) incubating a substrate-containing reaction medium with a cytochrome P450 monooxygenase derived from cytochrome P450 monooxygenase BM-3 from *Bacillus megaterium* having an amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43, 48-52.

CLAIMS AS FILED - 50915

67-70, 330-335, 352-356, 73-82 and 86-88; and

- b) isolating the oxidation product formed or a secondary product thereof from the medium;

where the monooxygenase mutant Phe87Val is not excluded.

- 14. A process as claimed in claim 13, wherein the exogenous or intermediately formed substrate is selected from:
 - a) optionally substituted mono- or polynuclear aromatics;
 - b) straight-chain or branched alkanes and alkenes;
 - c) optionally substituted cycloalkanes and cycloalkenes.
- 16. A process as claimed in claim 13, where the cytochrome P450 monooxygenase has at least one of the following mono- or polyamino acid substitutions:
 - a) Phe87Val;
 - b) Phe87Val, Leu188Gln; or
 - c) Phe87Val, Leu188Gln, Ala74Gly.
- 17. A process as claimed in claim 9, wherein, as exogenous substrate, at least one compound selected from the groups a) to d) of compounds defined above is added to a medium and the oxidation is carried out by enzymatic reaction of the substrate-containing medium in the presence of oxygen at a temperature of approximately 20 to 40°C and a pH of approximately 6 to 9, where the substrate-containing medium additionally contains an approximately 10- to 100-fold molar excess of reduction equivalents based on the substrate.

CLAIMS AS FILED - 50915

18. A process as claimed in claim 17, wherein, as exogenous substrate, a compound selected from indole, n-hexane, n-octane, n-decane, n-dodecane, cumene, 1-methylindole, α -, β - or γ -ionone, acridine, naphthalene, 6-methyl- or 8-methylquinoline, quinoline and quinaldine is employed.
19. A process for the microbiological production of indigo and/or indirubin, which comprises
 - a1) culturing a recombinant microorganism which produces an indole-oxidizing cytochrome P450 in a culture medium, in the presence of exogenous or intermediately formed indole; or
 - a2) incubating an indole-containing reaction medium with an indole-oxidizing cytochrome P450 monooxygenase; and
 - b) isolating the oxidation product formed or a secondary product thereof from the medium.
20. A process as claimed in claim 19, wherein the indigo and/or indirubin obtained, which was produced by oxidation of intermediately formed indole, is isolated from the medium.
21. A process as claimed in claim 20, wherein the indole oxidation is carried out by culturing the microorganisms in the presence of oxygen at a culturing temperature of approximately 20 to 40°C and a pH of approximately 6 to 9.
22. A process as claimed in claim 20, where the monooxygenase is derived from cytochrome P450 monooxygenase BM-3 from *Bacillus megaterium* having an

CLAIMS AS FILED - 50915

- amino acid sequence according to SEQ ID NO:2, which has at least one functional mutation in at least one of the amino acid sequence regions 172-224, 39-43, 48-52, 67-70, 330-335, 352-356, 73-82 and 86-88, including the substitution Phe87Val.
23. A process as claimed in claim 22, where the monooxygenase has at least one of the following mono- or polyamino acid substitutions:
- a) Phe87Val;
 - b) Phe87Val, Leu188Gln; or
 - c) Phe87Val, Leu188Gln, Ala74Gly.
24. A bioreactor comprising the cytochrom P450 monooxygenase as claimed in claim 1 or a recombinant microorganism transformed by a vector comprising an expression construct comprising a nucleic acid sequence coding for the cytochrom P450 monooxygenase of claim 1 in immobilized form.

Novel cytochrome P450 monooxygenases and their use for the
oxidation of organic compounds

- 5 The present invention relates to novel cytochrome P450 monooxygenases with modified substrate specificity which are capable of the oxidation of organic substrates, for example N-heterocyclic aromatic compounds, nucleotide sequences coding therefor, expression constructs and vectors comprising these
- 10 sequences, microorganisms transformed therewith, processes for the microbiological oxidation of various organic substrates, such as N-heterocyclic aromatic compounds and in particular processes for the preparation of indigo and indirubin.
- 15 Enzymes having novel functions and properties can be prepared either by screening of natural samples or by protein engineering of known enzymes. Under certain circumstances, the last-mentioned method can be the more suitable to induce characteristics whose generation by the natural selection route is improbable. Despite
- 20 numerous attempts at the engineering of enzymes, up to now there are only a few successful studies for promoting the catalytic activity of enzyme mutants with respect to a certain substrate (1-10). In these known cases, the substrates are structurally closely related to the native substrate of the respective enzyme.
- 25 As yet, there are no reports on the successful engineering of enzymes which, after modification, catalyze the reaction of a compound which structurally is completely different from the native substrate of the enzyme.
- 30 The cytochrome P450 monooxygenase isolatable from the bacterium *Bacillus megaterium* usually catalyzes the subterminal hydroxylation of long-chain, saturated acids and the corresponding amides and alcohols thereof or the epoxidation of unsaturated long-chain fatty acids or saturated fatty acids of
- 35 medium chain length (11-13). The optimal chain length of saturated fatty acids is 14 to 16 carbon atoms. Fatty acids having a chain length of less than 12 are not hydroxylated (11).

The structure of the heme domain of P450 BM-3 was determined by

40 X-ray structural analysis (14-16). The substrate binding site is present in the form of a long tunnel-like opening which extends from the surface of the molecule as far as the heme molecule and is almost exclusively bordered by hydrophobic amino acid residues. The only charged residues on the surface of the heme

45 domain are the residues Arg47 and Tyr51. It is assumed that these are involved in the binding of the carboxylate group of the substrate by formation of a hydrogen bond (14). The mutation of

2

- Arg47 to Glu brings about inactivation of the enzyme for arachidonic acid (13), but increases its activity compared with C₁₂-C₁₄-alkyltrimethylammonium compounds (17). Substrate utilization for aromatic compounds, in particular mono-, bi- or
- 5 polynuclear, if desired heterocyclic, aromatics, alkanes, alkenes, cycloalkanes and cycloalkenes, has not been described for this enzyme. Until now, it was therefore assumed in specialist circles that substrates other than the organic substrates hitherto described, for example indole, on account of
- 10 the clear structural differences from the native substrates of P450 BM-3, in particular on account of the absence of functional groups which could bind to the abovementioned residues in the substrate pocket, are not a substrate.
- 15 It is an object of the present invention to make available novel cytochrome P450 monooxygenases having modified substrate specificity or modified substrate profile. In particular, monooxygenase mutants are to be provided which, in comparison with the nonmutated wild-type enzyme, are enzymatically active
- 20 with structurally clearly different substrates.

Compared to the wild-type enzyme, a "modified substrate profile" can be observed for the mutants according to the invention. In particular, for the mutant in question, an improvement in

25 reactivity is observed, for example an increase of the specific activity (expressed as nmol of converted substrate/minute/nmol of P450 enzyme) and/or of at least one kinetic parameter selected from the group consisting of K_{cat}, K_m and K_{cat}/K_m (for example by at least 1%, such as 10 to 1000%, 10 to 500% or 10 to 100%) in

30 the conversion of at least one of the oxidizable compounds defined in groups a) to d). The oxidation reaction according to the invention comprises the enzyme-catalyzed oxygenation of at least one exogenous (i.e. added to the reaction medium) or endogenous (i.e. already present in the reaction medium) organic

35 substrate. In particular, the oxidation reaction according to the invention comprises a mono- and/or polyhydroxylation, for example a mono- and/or dihydroxylation, at an aliphatic or aromatic C-H group, or an epoxidation at a C=C group which is preferably non-aromatic. Also possible are combinations of the above

40 reactions. Moreover, the immediate reaction product can be converted further in the context of a non-enzymatic subsequent or side reaction. Such combinations of enzymatic and non-enzymatic processes likewise form part of the subject-matter of the present invention.

3

We have found that the above object is surprisingly achieved by means of novel cytochrome P450 monooxygenases which, for example, are capable of the oxidation of N-heterocyclic bi- or polynuclear aromatic compounds.

5

In particular, the invention relates to those monooxygenases whose substrate-binding region is capable by means of site-specific mutagenesis of the functional uptake of novel, for example N-heterocyclic substrates.

10

In a preferred embodiment of the invention, the novel monooxygenases are soluble, i.e. existent in non membrane-bound form, and enzymatically active in this form.

- 15 The monooxygenases according to the invention are preferably derived from cytochrome P450 monooxygenases of bacterial origin, as derived, in particular, from cytochrome P450 monooxygenase BM-3 from *Bacillus megaterium* having an amino acid sequence according to SEQ ID NO:2, which has at least one functional
- 20 mutation, i.e. promoting the oxidation of novel organic substrates (cf. in particular the groups a) to d) of compounds as defined below), for example N-heterocyclic mono-, bi- or polynuclear aromatic compounds, in one of the amino acid sequence regions 172-224 (F/G loop region), 39-43 (β -strand 1), 48-52
- 25 (β -strand 2), 67-70 (β -strand 3), 330-335 (β -strand 5), 352-356 (β -strand 8), 73-82 (helix 5) and 86-88 (helix 6).

The cytochrome P450 monooxygenase mutants provided according to the invention are preferably capable of at least one of the

30 following reactions:

- a) oxidation of unsubstituted or substituted N-, O- or S-heterocyclic mono-, bi- or polynuclear aromatic compounds;
- b) oxidation of unsubstituted or substituted mono- or
- 35 polynuclear aromatics;
- c) oxidation of straight-chain or branched alkanes and alkenes; and
- d) oxidation of unsubstituted or substituted cycloalkanes and cycloalkenes.

40

Preferred monooxygenase mutants have at least one functional mutation, in particular amino acid substitution, in at least one of the sequence regions 73-82, 86-88 and 172-224. Thus, for example, Phe87 can be replaced by an amino acid having an

45 aliphatic side chain, such as Ala, Val, Leu, in particular Val; Leu188 can be replaced by an amino acid having an amide side chain, such as Asn or, in particular, Gln; and Ala74 can be

0050/50915

4

replaced by another amino acid having an aliphatic side chain, such as Val and, in particular, Gly.

Particularly preferred monooxygenase mutants of this type are those which have at least one of the following mono- or polyamino acid substitutions:

- 1) Phe87Val;
- 2) Phe87Val, Leu188Gln; or
- 10 3) Phe87Val, Leu188Gln, Ala74Gly;

and functional equivalents thereof. The number indicates the position of the mutation; the original amino acid is indicated before the number and the newly introduced amino acid after the number.

In this context, "functional equivalents" or analogs of the mutants which are disclosed specifically are mutants differing therefrom which furthermore have the desired substrate specificity with respect to at least one of the oxidation reactions a) to d) described above, i.e., for example, for heterocyclic aromatics and which hydroxylate, for example, indole, or furthermore exhibit the desired "modified substrate profile" with respect to the wild-type enzyme.

"Functional equivalents" are also to be understood as meaning in accordance with the invention mutants which exhibit, in at least one of the abovementioned sequence positions, an amino acid substitution other than the one mentioned specifically, but still lead to a mutant which, like the mutant which has been mentioned specifically, show a "modified substrate profile" with respect to the wild-type enzyme and catalyze at least one of the abovementioned oxidation reactions. Functional equivalence exists in particular also in the case where the modifications in the substrate profile correspond qualitatively, i.e. where, for example, the same substrates are converted, but at different rates.

"Functional equivalents" naturally also encompass P450 monooxygenase mutants which, like the P450 BM3 mutants which have been mentioned specifically, can be obtained by mutating P450 enzymes from other organisms. For example, regions of homologous sequence regions can be identified by sequence comparison. Following the principles of what has been set out specifically in the invention, the modern methods of molecular modeling then

0050/50915

5

allow equivalent mutations to be carried out which affect the reaction pattern.

"Functional equivalents" also encompass the mutants which can be
5 obtained by one or more additional amino acid additions, substitutions, deletions and/or inversions, it being possible for the abovementioned additional modifications to occur in any sequence position as long as they give rise to a mutant with a modified substrate profile in the above sense.

10

Substrates of group a) which can be oxidized according to the invention are unsubstituted or substituted heterocyclic mono-, bi- or polynuclear aromatic compounds; in particular oxidizable or hydroxylatable N-, O- or S-heterocyclic mono-, bi- or
15 polynuclear aromatic compounds. They include preferably two or three, in particular two, 4- to 7-membered, in particular 6- or 5-membered, fused rings, where at least one, preferably all, rings have aromatic character and where at least one of the aromatic rings carries one to three, preferably one, N-, O- or
20 S-heteroatom in the ring. The total ring structure may contain one or two further identical or different heteroatoms. The aromatic compounds may furthermore carry 1 to 5 substituents at the ring carbon or heteroatoms. Examples of suitable substituents are C₁- to C₄-alkyl, such as methyl, ethyl, n- or isopropyl, n-,
25 iso- or t-butyl, or C₂- to C₄-alkenyl, such as ethenyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl or 3-butenyl, hydroxyl and halogen, such as F, Cl and Br. The alkyl or alkenyl substituents mentioned may also have a keto or aldehyde group; examples being propan-2-on-3-yl, butan-2-on-4-yl,
30 3-buten-2-on-4-yl. Non-limiting examples of suitable heterocyclic substrates are, in particular, binuclear heterocycles, such as indole, N-methyl-indole, and the substituted analogs thereof which carry one to three of the above-defined substituents on carbon atoms, for example 5-chloro- or 5-bromoindole; and also
35 quinoline and quinoline derivatives, for example 8-methylquinoline, 6-methyl-quinoline and quinaldine; and benzothiophene, and the substituted analogs thereof which carry one to three of the above-defined substituents on carbon atoms. Moreover, trinuclear hetero-aromatics, such as acridine and the
40 substituted analogs thereof which carry one to three of the above-defined substituents on carbon atoms, may be mentioned.

Substrates of group b) which are oxidizable according to the invention are unsubstituted or substituted mono- or polynuclear,
45 in particular mono- or binuclear, aromatics, such as benzene and naphthalene. The aromatic compounds may be unsubstituted or mono- or polysubstituted and, for example, carry 1 to 5 substituents on

0050/50915

6

the ring carbon atoms. Examples of suitable substituents are C₁- to C₄-alkyl, such as methyl, ethyl, n- or isopropyl or n-, iso- or t-butyl, or C₂- to C₄-alkenyl, such as ethenyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl or 3-butenyl, hydroxyl and
5 halogen, such as F, Cl and Br. The alkyl or alkenyl substituents mentioned may also have a keto or aldehyde group; Examples being propan-2-on-3-yl, butan-2-on-4-yl, 3-buten-2-on-4-yl. The aromatic may be fused with a four- to seven-membered non-aromatic ring. The non-aromatic ring may have one or two C=C double bonds,
10 be mono- or polysubstituted by the abovementioned substituents and may carry one or two hetero ring atoms. Examples of particularly suitable aromatics are mononuclear aromatics, such as cumene, and binuclear substrates, such as indene and naphthalene, and substituted analogs thereof which carry one to
15 three of the above-defined substituents on carbon atoms.

Substrates of group c) which can be oxidized according to the invention are straight-chain or branched alkanes or alkenes having 4 to 15, preferably 6 to 12, carbon atoms. Examples which
20 may be mentioned are n-butane, n-pentane, n-hexane, n-heptane, n-octane, n-nonane, n-decane, n-undecane and n-dodecane, and the analogs of these compounds which are branched once or more than once, for example analogous compounds having 1 to 3 methyl side groups; or mono- or polyunsaturated, for example
25 mono-unsaturated, analogs of the abovementioned alkanes.

Substrates of group d) which can be oxidized according to the invention are unsubstituted or substituted cycloalkanes and cycloalkenes having 4 to 8 ring carbon atoms. Examples of these
30 are cyclopentane, cyclopentene, cyclohexane, cyclohexene, cycloheptane and cycloheptene. The ring structure may carry one or more, for example 1 to 5, substituents according to the above definition for compounds of groups a) and b). Nonlimiting examples are ionones, such as α -, β - and γ -ionone, and the
35 corresponding methyl ionones and iso-methyl ionones. Particular preference is given to α - and β -ionone.

The invention also relates to nucleic acid sequences coding for one of the monooxygenases according to the invention. Preferred
40 nucleic acid sequences are derived from SEQ ID NO:1, which have at least one nucleotide substitution which leads to one of the functional amino acid mutations described above. The invention moreover relates to functional analogs of the nucleic acids obtained by addition, substitution, insertion and/or deletion of
45 individual or multiple nucleotides, which furthermore code for a

7

monooxygenase having the desired substrate specificity, for example having indole-oxidizing activity.

- The invention also encompasses those nucleic acid sequences which
- 5 comprise so-called silent mutations or which are modified in comparison with a specifically mentioned sequence in accordance with the codon usage of a specific origin or host organism, and naturally occurring variants of such nucleic acid sequences. The invention also encompasses modifications of the nucleic acid
- 10 sequences obtained by degeneration of the genetic code (i.e. without any changes in the corresponding amino acid sequence) or conservative nucleotide substitution (i.e. the corresponding amino acid is replaced by another amino acid of the same charge, size, polarity and/or solubility), and sequences modified by
- 15 nucleotide addition, insertion, inversion or deletion, which sequences encode a monooxygenase according to the invention having a "modified substrate profile", and the corresponding complementary sequences.
- 20 The invention furthermore relates to expression constructs comprising a nucleic acid sequence encoding a mutant according to the invention under the genetic control of regulatory nucleic acid sequences; and vectors comprising at least one of these expression constructs.
- 25 Preferably, the constructs according to the invention encompass a promoter 5'-upstream of the encoding sequence in question and a terminator sequence 3'-downstream, and, optionally, further customary regulatory elements, and, in each case operatively
- 30 linked with the encoding sequence. Operative linkage is to be understood as meaning the sequential arrangement of promoter, encoding sequence, terminator and, if appropriate, other regulatory elements in such a manner that each of the regulatory elements can fulfill its intended function on expression of the
- 35 encoding sequence. Examples of operatively linkable sequences are targeting sequences, or else translation enhancers, enhancers, polyadenylation signals and the like. Further regulatory elements encompass selectable markers, amplification signals, replication origins and the like.
- 40 In addition to the artificial regulatory sequences, the natural regulatory sequence can still be present upstream of the actual structural gene. If desired, this natural regulation may be switched off by genetic modification, and the expression of the
- 45 genes may be enhanced or lowered. However, the gene construct may also be simpler in construction, i.e. no additional regulatory signals are inserted upstream of the structural gene and the

0050/50915

8

natural promoter with its regulation is not removed. Instead, the natural regulatory sequence is mutated in such a way that regulation no longer takes place and the gene expression is increased or reduced. One or more copies of the nucleic acid sequences may be present in the gene construct.

Examples of suitable promoters are: cos, tac, trp, tet, trp-tet, lpp, lac, lpp-lac, lacIq, T7, T5, T3, gal, trc, ara, SP6, l-PR or l-PL promoter, which are advantageously employed in Gram-negative bacteria; and Gram-positive promoters amy and SPO2, the yeast promoters ADC1, MFA, Ac, P-60, CYC1, GAPDH or the plant promoters CaMV/35S, SSU, OCS, lib4, usp, STLS1, B33, nos or the ubiquitin or phaseolin promoter. Particular preference is given to using inducible promoters, for example light- and in particular temperature-inducible promoters, such as the P_{rP1} promoter.

In principle, all natural promoters with their regulatory sequences can be used. In addition, synthetic promoters may also be used in an advantageous fashion.

The abovementioned regulatory sequences are intended to allow the targeted expression of the nucleic acid sequences and of protein expression. Depending on the host organism, this may mean, for example, that the gene is expressed or overexpressed only after induction has taken place, or that it is expressed and/or overexpressed immediately.

The regulatory sequences or factors can preferably have a positive effect on expression and in this manner increase or reduce the latter. Thus, an enhancement of the regulatory elements may advantageously take place at the transcriptional level by using strong transcription signals such as promoters and/or "enhancers". In addition, translation may also be enhanced by improving, for example, mRNA stability.

An expression cassette is generated by fusing a suitable promoter with a suitable monooxygenase nucleotide sequence and a terminator signal or polyadenylation signal. To this end, customary recombination and cloning techniques are used as they are described, for example, in T. Maniatis, E.F. Fritsch and J. Sambrook, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989) and in T.J. Silhavy, M.L. Berman and L.W. Enquist, Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1984) and in Ausubel, F.M. et al., Current Protocols in

Molecular Biology, Greene Publishing Assoc. and Wiley Interscience (1987).

For expression in a suitable host organism, the recombinant
5 nucleic acid construct or gene construct is advantageously
inserted into a host-specific vector which allows optimal gene
expression in the host. Vectors are well known to the skilled
worker and can be found, for example, in "Cloning Vectors"
(Pouwels P.H. et al., Ed., Elsevier, Amsterdam-New York-Oxford,
10 1985). Vectors are to be understood as meaning not only plasmids,
but all other vectors known to the skilled worker such as, for
example, phages, viruses, such as SV40, CMV, baculovirus and
adenovirus, transposons, IS elements, phasmids, cosmids, and
linear or circular DNA. These vectors can be replicated
15 autonomously in the host organism or chromosomally.

The vectors according to the invention allow the generation of
recombinant microorganisms which are transformed, for example,
with at least one vector according to the invention and which can
20 be employed for producing the mutants. The above-described
recombinant constructs according to the invention are
advantageously introduced into a suitable host system and
expressed. It is preferred to use usual cloning and transfection
methods known to the skilled worker in order to bring about
25 expression of the abovementioned nucleic acids in the expression
system in question. Suitable systems are described, for example,
in current protocols in molecular biology, F. Ausubel et al.,
Ed., Wiley Interscience, New York 1997.

30 Suitable host organisms are, in principle, all organisms which
allow expression of the nucleic acids according to the invention,
their allelic variants, and their functional equivalents or
derivatives. Host organisms are to be understood as meaning, for
example, bacteria, fungi, yeasts or plant or animal cells.
35 Preferred organisms are bacteria such as those of the genera
Escherichia, such as, for example, Escherichia coli,
Streptomyces, Bacillus or Pseudomonas, eukaryotic microorganisms
such as Saccharomyces cerevisiae, Aspergillus, and higher
eukaryotic cells from animals or plants, for example Sf9 or CHO
40 cells.

If desired, expression of the gene product may also be brought
about in transgenic organisms such as transgenic animals such as,
in particular, mice, sheep, or transgenic plants. The transgenic
45 organisms may also be knock-out animals or plants in which the

0050/50915

10

corresponding endogenous gene has been eliminated, such as, for example, by mutation or partial or complete deletion.

Successfully transformed organisms can be selected by marker
5 genes which are likewise contained in the vector or in the
expression cassette. Examples of such marker genes are genes for
resistance to antibiotics and for enzymes which catalyze a color
reaction, which causes staining of the transformed cell. These
transformed cells can then be selected using automatic cell
10 selection. Microorganisms which have been transformed
successfully with a vector and which carry an appropriate gene
for resistance to antibiotics (for example G418 or hygromycin)
can be selected by using appropriate antibiotics-containing media
or substrates. Marker proteins which are presented on the cell
15 surface can be used for selection by affinity chromatography.

The combination of the host organisms and the vectors appropriate
for the organisms, such as plasmids, viruses or phages, such as,
for example, plasmids with the RNA polymerase/promoter system,
20 phages λ , μ or other temperate phages or transposons and/or other
advantageous regulatory sequences forms an expression system. The
term "expression system" means, for example, a combination of
mammalian cells such as CHO cells, and vectors, such as pcDNA3neo
vector, which are suitable for mammalian cells.

25 As described above, the gene product can also be expressed
advantageously in transgenic animals, for example mice, sheep, or
transgenic plants. It is likewise possible to program cell-free
translation systems with the RNA derived from the nucleic acid.

30 The invention furthermore provides a process for preparing a
monooxygenase according to the invention, which comprises
cultivating a monooxygenase-producing microorganism, if
appropriate inducing the expression of the monooxygenase, and
35 isolating the monooxygenase from the culture. If desired, the
monooxygenase according to the invention can thus also be
produced on an industrial scale.

The microorganism can be cultivated and fermented by known
40 methods. Bacteria, for example, can be grown in a TB or LB medium
and at 20-40°C and a pH of 6-9. Suitable cultivation conditions
are described in detail in T. Maniatis, E.F. Fritsch and
J. Sambrook, Molecular Cloning: A Laboratory Manual, Cold Spring
Harbor Laboratory, Cold Spring Harbor, NY (1989), for example.

45

11

- If the monooxygenase is not secreted into the culture medium, the cells are then lyzed and the monooxygenase is obtained from the lysate using known methods for the isolation of proteins. The cells can be lyzed alternatively by high-frequency ultrasound, by high pressure, for example in a French pressure cell, by osmolysis, by the action of detergents, lytic enzymes or organic solvents, by homogenization or by a combination of a plurality of the processes mentioned. Purification of the monooxygenase can be achieved by known chromatographic processes, such as molecular sieve chromatography (gel filtration), such as Q-Sepharose chromatography, ion-exchange chromatography and hydrophobic chromatography, and by other customary processes, such as ultrafiltration, crystallization, salting out, dialysis and native gel electrophoresis. Suitable processes are described, for example, in Cooper, F.G., Biochemische Arbeitsmethoden [Biochemical Procedures], Verlag Walter de Gruyter, Berlin, New York or in Scopes, R., Protein Purification, Springer Verlag, New York, Heidelberg, Berlin.
- To isolate the recombinant protein, it is particularly advantageous to use vector systems or oligonucleotides which extend the cDNA by certain nucleotide sequences and thus code for modified polypeptides or fusion proteins which serve to simplify purification. Suitable modifications of this type are, for example, so-called "tags" which act as anchors, such as, for example, the modification known as hexa-histidine anchor, or epitopes which can be recognized as antigens by antibodies (described, for example, in Harlow, E. and Lane, D., 1988, Antibodies: A Laboratory Manual. Cold Spring Harbor (N.Y.) Press). These anchors can be used to attach the proteins to a solid support such as, for example, a polymer matrix, which can, for example, be packed into a chromatography column, or to a microtiter plate or to another support.
- These anchors can also at the same time be used to recognize the proteins. It is also possible to use for recognition of the proteins conventional markers such as fluorescent dyes, enzyme markers which form a detectable reaction product after reaction with a substrate, or radioactive markers, alone or in combination with the anchors for derivatizing the proteins.

The invention moreover relates to a process for the microbiological oxidation of organic compounds, for example N-heterocyclic mono-, bi- or polynuclear aromatic compounds according to the above definition, which comprises

12

- a1) culturing a recombinant microorganism according to the above definition in a culture medium, in the presence of an exogenous (added) substrate or an intermediately formed substrate, which substrate is oxidizable by the monooxygenase according to the invention, preferably in the presence of oxygen (i.e. aerobically); or
- a2) incubating a substrate-containing reaction medium with an enzyme according to the invention, preferably in the presence of oxygen and an electron donor; and
- 10 b) isolating the oxidation product formed or a secondary product thereof from the medium.

The oxygen required for the reaction either passes from the atmosphere into the reaction medium or, if required, can be added in a manner known per se.

The oxidizable substrate is preferably selected from

- a) unsubstituted or substituted N-heterocyclic mono-, bi- or polynuclear aromatic compounds;
- b) unsubstituted or substituted mono- or polynuclear aromatics;
- c) straight-chain or branched alkanes and alkenes;
- d) unsubstituted or substituted cycloalkanes and cycloalkenes.

25 A preferred process variant is directed to the formation of indigo/indirubin and is characterized by the fact that the substrate is indole formed as an intermediate in the culture and that the indigo and/or indirubin formed in the culture medium is isolated by oxidation of hydroxyindole intermediates.

30

If the oxidation according to the invention is carried out using a recombinant microorganism, the culturing of the microorganisms is preferably first carried out in the presence of oxygen and in a complex medium, such as, for example, TB or LB medium at a culturing temperature of approximately 20 to 40°C and a pH of approximately 6 to 9, until an adequate cell density is reached. The addition of exogenous indole is usually not necessary, as this is intermediately formed by the microorganism. However, when using other substrates, addition of exogenous substrate may be required. In order to be able to control the oxidation reaction better, the use of an inducible, in particular temperature-inducible, promoter is preferred. The temperature is in this case increased to the necessary induction temperature, e.g. 42°C in the case of the P_{rP_1} promoter, this is maintained for a sufficient period of time, e.g. 1 to 10 or 5 to 6 hours, for the expression of the monooxygenase activity and the temperature is then reduced again to a value of approximately 30 to 40°C. The culturing is


0050/50915

13

then continued in the presence of oxygen for 12 hours to 3 days. The pH can, in particular in the case of indole oxidation, be increased by addition of NaOH, e.g. to 9 to 10, whereby the indigo formation or indirubin formation is additionally promoted
 5 by atmospheric oxidation of the enzymatically formed oxidation products 2- and 3-hydroxyindole.

The indigo/indirubin formation according to the invention is illustrated by the reaction scheme below:

10



45 However, if the oxidation according to the invention is carried out using purified or enriched enzyme mutants, the enzyme according to the invention is dissolved in an exogenous

0050/50915

14

substrate-containing, for example indole-containing medium (approximately 0.01 to 10 mM, or 0.05 to 5 mM), and the reaction is carried out, preferably in the presence of oxygen, at a temperature of approximately 10 to 50°C, such as, for example, 30
5 to 40°C, and a pH of approximately 6 to 9 (such as established, for example, using 100 to 200 mM phosphate or tris buffer), and in the presence of a reductant, the substrate-containing medium moreover containing, relative to the substrate to be oxidized, an approximately 1- to 100-fold or 10- to 100-fold molar excess of
10 reduction equivalents. The preferred reductant is NADPH. If required, the reducing agent can be added in portions.

In a similar manner, the oxidizable substrates which are preferably used are: n-hexane, n-octane, n-decane, n-dodecane,
15 cumene, 1-methylindole, 5-Cl- or Br-indole, indene, benzothiophene, α -, β - and γ -ionone, acridine, naphthalene, 6-methyl- or 8-methylquinoline, quinoline and quinaldine.

The enzymatic oxidation reaction according to the invention can
20 be carried out, for example, under the following conditions:

Substrate concentration:	from 0.01 to 20 mM
Enzyme concentration:	from 0.1 to 10 mg/ml
25 Reaction temperature:	from 10 to 50°C
pH:	from 6 to 8
30 Buffer:	from 0.05 to 0.2 M potassium phosphate, or Tris/HCl
Electron donor:	is preferably added in portions (initial concentration about 35 0.1 to 2 mg/ml)

The mixture can briefly (from 1 to 5 minutes) be preincubated (at about 20-40°C) before the reaction is initiated, for example by adding the electron donors (e.g. NADPH). The reaction is carried
40 out aerobically, if appropriate with additional introduction of oxygen.

In the substrate oxidation process according to the invention, oxygen which is present in or added to the reaction medium is
45 cleaved reductively by the enzyme. The required reduction

0050/50915

15

equivalents are provided by the added reducing agent (electron donor).

The oxidation product formed can then be separated off from the
5 medium and purified in a conventional manner, such as, for example, by extraction or chromatography.

Further subjects of the invention relate to bioreactors, comprising an enzyme according to the invention or a recombinant
10 microorganism according to the invention in immobilized form.

A last subject of the invention relates to the use of a cytochrome P450 monooxygenase according to the invention or of a vector or microorganism according to the invention for the
15 microbiological oxidation of a substrate from one of the groups a) to d), in particular of N-heterocyclic mono-, bi- or polynuclear aromatic compounds, and preferably for the formation of indigo and/or indirubin.

20 The present invention is now described in greater detail with reference to the following examples.

Example 1:

25 Randomization of specific codons of P450 BM-3

The experiments were carried out essentially as described in (19). Three positions (Phe87, Leu188 and Ala74) were randomized with the aid of site-specific mutagenesis using the Stratagene
30 QuikChange kit (La Jolla, CA, USA). The following PCR primers were used for the individual positions:

Phe87: 5'-gcaggagacgggttgnnnacaagctggacg-3' (SEQ ID NO:3),
5'-cgtccagcttgtnnncaacccgtctcctgc-3', (SEQ ID NO:4)
35 Leu188: 5'-gaagcaatgaacaagnnnacagcgagcaaattccag-3' (SEQ ID NO:5),
5'-ctggatttgctcgctgtnnncttgcttcattgcttc-3' (SEQ ID NO:6);
Ala74: 5'-gctttgataaaaacttaaagtcaannncttaaatttgtagc-3' (SEQ ID:
NO:7),
5'-cgtacaaatttaagnnncttgacttaagtttttatcaaagc-3' (SEQ ID
40 NO:8)

The conditions for the PCR were identical for all three positions. In particular, 17.5 pmol of one of each primer, 20 pmol of template plasmid DNA, 3 U of the Pfu polymerase and
45 3.25 nmol of each dNTP were used per 50 µl reaction volume. The PCR reaction was started at 94°C/1 min and the following temperature cycle was then carried out 20 times: 94°C, 1 min;

0050/50915

16

46°C, 2.5 min; 72°C, 17 min. After 20 cycles, the reaction was continued at 72°C for 15 min. After the PCR, the template DNA was digested at 37°C for 3 h using 20 U of DpnI. *E. coli* DH5 α was then transformed. The transformed *E. coli* DH5 α cells were plated out
5 onto LB agar plates which contained 150 μ g/ml of ampicillin. Incubation was then carried out at 37°C for 18 h.

Example 2:

Expression and purification of the P450 BM-3 and its mutants and
10 production of a blue pigment

The P450 BM-3 gene and the mutants thereof were expressed under the control of the strong, temperature-inducible P_{RP_L} promoter of the plasmid pCYTEXP1 in *E. coli* DH5 α as already described (20).
15 Colonies were picked up using sterile toothpicks and transferred to microtiter plates having 96 hollows, comprising 200 μ l of TB medium and 100 μ g/ml of ampicillin per hollow. Incubation was then carried out at 37°C overnight. 40 μ l of the cell culture of one of each hollow were then transferred to a culture tube which
20 contained 2 ml of TB medium with 100 μ g/ml of ampicillin. Culturing was then carried out at 37°C for 2 h. The temperature was then increased to 42°C for 6 h for induction. Culturing was then continued at 37°C overnight, a blue pigment being produced.

25 The preparative production of enzyme or blue pigment was carried out starting from a 300 ml cell culture ($OD_{578nm} = 0.8$ to 1.0). For the isolation of the enzyme, the cells were centrifuged off at 4000 rpm for 10 min and resuspended in 0.1 M K_xPO₄ buffer, pH 7.4. The ice-cooled cells were carefully disrupted with the aid
30 of a Branson sonifer W25 (Dietzenbach, Germany) at an energy output of 80 W by 2 min sonification three times. The suspensions were centrifuged at 32570 x g for 20 min. The crude extract was employed for the activity determination or for the enzyme purification. The enzyme purification was carried out as already
35 described in (21), to which reference is expressly made hereby. The concentration of purified enzyme was determined by means of the extinction difference at 450 and 490 nm, as already described in (11), using an extinction coefficient ϵ of 91 mM⁻¹ cm⁻¹.

40 Example 3:

Isolation of mutants which produce large amounts of blue pigment

100 colonies in each case were isolated from the mutants of one
45 of each position, which were produced by randomized mutagenesis of the codon of the corresponding position. These colonies were cultured in culture tubes for the production of blue pigment.

0050/50915

17

After washing the cells with water and a number of slow centrifugation steps (500 rpm), the blue pigment was extracted using dimethyl sulfoxide (DMSO). The solubility of the blue pigment was greatest in DMSO. The absorption of the extract was
5 determined at 677 nm. That mutant which produced the largest amount of blue pigment, especially mutants from a specific position, was used for DNA sequencing (ABI DNA sequencing kit; ABI Prism™ 377 DNA sequencer) and moreover as a template for site-specific randomized mutagenesis.

10

Example 4:

Activity test for the indole hydroxylation

15 The indole hydroxylation activity was tested in a solution which contained 8 μ l of a 10-500 mM indole solution in DMSO, 850 μ l of tris/HCl buffer (0.1 M, pH 8.2) and 0.6 nmol of P450 BM-3 wild type or mutant in a final volume of 1 ml. The mixture was preincubated for 9 min before the reaction was started by
20 addition of 50 μ l of an aqueous 1 mM solution of NADPH. The reaction was stopped after 20 sec by addition of 60 μ l of 1.2 M KOH. Within 5 to 30 sec (under aerobic conditions), the enzyme products were converted completely into indigo [$\Delta^{2,2'}$ -biindoline]-3,3'-dione) and indirubin
25 ([$\Delta^{2,3'}$ -biindoline]-2',3-dione). The indigo production was determined by means of its absorption at 670 nm. A calibration curve using pure indigo showed an extinction coefficient of 3.9 mM⁻¹ cm⁻¹ at this wavelength. A linear curve was obtained for indigo production in a reaction time of 40 sec using 0.6 nmol of
30 wild type or P450 BM-3 mutant and 0.05 to 5.0 mM of indole. Indirubin shows a very weak absorption at 670 nm and the amount of indirubin formed was very much smaller than the amount of indigo formed. The formation of indirubin was neglected in the determination of the kinetic parameters. The NADPH consumption
35 was determined at 340 nm and calculated as described (17) using an extinction coefficient of 6.2 mM⁻¹ cm⁻¹.

Example 5:

40 Purification of indigo and indirubin

After washing the cells with water and repeated centrifugation at 500 g, the blue pellet formed was extracted using tetrahydrofuran (THF). The extract was evaporated almost to dryness and the red
45 pigment was extracted a number of times with 50 ml of absolute ethanol. The residual blue solid was dissolved in THF and analyzed by thin-layer chromatography (TLC). The ethanol solution

0050/50915

18

was evaporated and purified by silica gel chromatography (TLC 60, Merck, Darmstadt, Germany; 2 cm x 30 cm) before it was washed with THF and petroleum ether in a ratio of 1:2. The red solution obtained was evaporated and its purity was determined by TLC. The
5 absorption spectra of the blue and of the red pigment were determined in a range from 400 to 800 nm with the aid of an Ultraspec 3000 spectrophotometer (Pharmacia, Uppsala, Sweden). The blue and the red color were moreover analyzed by mass spectrometry and ¹H-NMR spectroscopy.

10

Experimental results

1. Increasing the productivity for blue pigment by P450 BM-3 mutagenesis

15

Native P450 BM-3 does not have the ability to produce the blue indigo-containing pigment, or the precursor substances 2- or 3-hydroxyindole. In order to be able to prepare a sufficient amount of blue pigment, P450 BM-3 was subjected to evolution in a
20 controlled manner. All mutants which produced the blue pigment were sequenced. It was found that at least one of the following three positions were mutated: Phe87, Leu188 and Ala74. It was therefore assumed that these three positions play a crucial role for the activity of P450 BM-3 in the production of blue pigment.
25 From the structure of the heme domain of cytochrome P450 BM-3, complexed with palmitoleic acid, it is seen that Phe87 prevents the substrate from coming closer to the heme group (14). The mutant Phe87Val shows a high regio- and stereoselectivity in the epoxidation of (14S, 15R)-arachidonic acid (13) and the mutant
30 Phe87Ala shifts the hydroxylation position of ω -1, ω -2 and ω -3 to ω (22). The position 87 was therefore selected as first for the site-specific randomized mutagenesis with the aid of PCR. In tube cultures, 7 colonies were obtained which produced a small amount of blue pigment after induction. The colony which produced the
35 largest amount of the blue pigment was selected for the DNA sequencing. The sequence data showed substitution of Phe87 by Val. The mutant Phe87Val was then used as a template for the second round of site-specific randomized mutagenesis on position Leu188. The structure of the heme domain, complexed with
40 palmitoleic acid, shows that the repositioning of the F and G helices brings the residue Leu188 into direct contact with the substrate (14). This position can therefore play an important role in substrate binding or orientation. After the second screening passage, 31 colonies were observed which produced the
45 blue pigment. The mutant which produced the largest amount of pigment contained the substitutions Phe87Val and Leu188Gln. This mutant was then mutated in position Ala74 in the third passage of

0050/50915

19

site-specific randomized mutagenesis. In this case the triple mutant F87L188A74 (Phe87Val, Leu188Gln and Ala74Gly) was obtained, which produced several mg of blue pigment in a 2-liter flask, containing 300 ml of TB medium. This amount was sufficient for the isolation and characterization of the blue pigment.

2. Isolation and identification of the blue pigment

After washing the cells, the residual blue pellet was extracted with THF and analyzed by TLC. The blue pigment was separated into a rapidly migrating blue component and into a more slowly migrating red component. Both components showed exactly the same mobility parameters as the components of a commercial indigo sample.

After the purification, the absorption spectra of both components were determined in DMSO. The blue component showed the same spectrum as a commercial indigo sample. The purified blue and red components were each analyzed by mass spectrometry. The mass spectra of both pigments showed a strong molecular ion peak at $m/e = 262$ and two fragment peaks at $m/e = 234$ and 205 (relative intensity in each case 10%). This pattern is typical of indigoid compounds. The elementary composition of these ions was determined by high-resolution mass spectrometry as $C_{16}H_{10}N_2O_2$, $C_{15}H_{10}N_2O$ and $C_{14}H_9N_2$. This is also characteristic of structures of the indigo type. The blue pigment was thus identified as indigo and the red pigment as indirubin. For the confirmation of the structure, 500 MHz 1H -NMR spectra of both pigments were carried out in DMSO- D_6 solution. The results agreed with the literature data (23).

3. Production of indigo using isolated enzymes

It is known that indigo is accessible from indole by microbial transformation (24-26). None of these microbial systems, however, contained a P450 monooxygenase. According to the invention, the catalytic activity of the pure enzyme for indole was first determined. The mutant F87L188A74 was mixed with indole. No color reaction could be observed. Only after addition of NADPH to the reaction mixture was the blue pigment formed after approximately 20 min. By adjustment of the pH of the reaction mixture to a value of approximately 11, 30 sec after addition of NADPH, the blue coloration was visible within a few seconds. Control experiments using native P450 BM-3 were always negative, even using increased concentrations of enzyme, indole and NADPH. The blue pigment was extracted using ethyl acetate and analyzed by TLC. The blue pigment again separated into a more rapidly running

0050/50915

20

blue component and into a slower running red component. The R_f values and the absorption spectra were identical to those values of the extracts from the fermentation broth. The F87L188A74 mutant of P450 BM-3 is thus an indole hydroxylase.

5

Two routes have previously been described for the enzymatic transformation of indole to indigo. One route is catalyzed by a dioxygenase, the other by a styrene monooxygenase (24, 25). The NADPH stoichiometry is in both cases 2. It was therefore assumed

10 that in contrast to the dioxygenases the mutant F87L188A74 according to the invention hydroxylates indole in only one position to form oxindole (2-hydroxyindole) or indoxyl (3-hydroxyindole).

15 4. Kinetic parameters of indole hydroxylation

Pure samples of the wild-type enzyme P450 BM-3 and of the mutants Leu188Gln, Phe87Val, F87L188 and F87L188A74 were used for the determination of the kinetic parameters of indole hydroxylation.

20 The results are summarized in Table 1 below.

Table 1: Kinetic parameters of the P450 BM-3 mutants for indole hydroxylation

25 Mutants	K_{cat} (S^{-1})	K_m (mM)	K_{cat}/K_m ($M^{-1}s^{-1}$)
WT	-a)	-	-
Leu188Gln	n.d. ^{b)}	n.d.	n.d.
Phe87Val	2.03 (0.14)	17.0 (1.0)	119
F87L188	2.28 (0.16)	4.2 (0.4)	543
30 F87L188A74	2.73 (0.16)	2.0 (0.2)	1365

a) no activity was observed;

b) not determined (activity was too low to be measured)

35 Even with an excess of purified enzyme and high indole concentration, the wild-type enzyme is not able to oxidize indole. The mutant Leu188Gln shows a low activity. The mutant Phe87Val shows a catalytic activity of $119 M^{-1}s^{-1}$ for indole hydroxylation. The catalytic efficiency of the double mutant F87L188 (Phe87Val, Leu188Gln) increased to $543 M^{-1}s^{-1}$ and was

40 increased to $1365 M^{-1}s^{-1}$ by introduction of the further substitution Ala74Gly. The K_{cat} values increased from Phe87Val to the triple mutant by a total of 35%, while the K_m values decreased approximately by seven-fold. This indicates that Ala74Gly and Leu188Gln are mainly involved in substrate binding.

45

0050/50915

21

For the triple mutant F87L188A74, the indole turnover rate ($K_{cat}=2.73 \text{ s}^{-1}$) was more than ten times higher than for most P450 enzymes (18).

5 Example 6

Hydroxylation of n-octane using modified cytochrome P450 monooxygenase

- 10 The reactions were carried out using a P450 BM-3 monooxygenase mutant comprising the following mutations: Phe87Val Leu188Gln Ala74Gly

The chosen substrate was n-octane. For the hydroxylation of
15 n-octane, the following aerobic reaction mixture was used:

- | | |
|-------------------|---|
| P450 BM-3 mutant: | 17.5 mg (lyophilisate) |
| Reaction buffer: | 9.1 ml (potassium phosphate buffer 50 mM, pH 7.5) |
| 20 Substrate: | 50 μl of a 60 mM solution (in acetone) |
| Temperature: | 25°C |

- The enzyme lyophilisate was dissolved in 500 μl of reaction buffer and initially incubated at room temperature with substrate and
25 reaction buffer for 5 minutes. 300 μl NADPH solution (5 mg/ml) were then added. Addition of NADPH was repeated two more times. The progress of the reaction was monitored by measuring the absorption at 340 nm, which allows the NADPH decrease to be observed. NADPH is added in aliquots of 300 μl , since too high a
30 concentration of NADPH in the reaction solution would result in inactivation of the enzyme. To isolate the products, the reaction solution was then extracted three times with 5 ml of diethyl ether. The combined organic phases were dried over MgSO_4 and concentrated. The products were then characterized by TLC, GC/MS
35 and NMR.

The GC/MS analysis of the reaction mixture gave the following result:

40 Compound	Rt[min] ¹⁾	Conversion [%]
4-octanol	13.51	37
3-octanol	14.08	47
2-octanol	14.26	16

- 45 1) Temperature program: 40°C 1 min isothermic / 3°C/min 95°C /10°C/min 275°C; apparatus: Finnigan MAT 95; GC: HP 5890 Series II

0050/50915

22

Split Injector; Column: HP-5MS (methylsiloxane) 30m x 0.25mm;
Carrier gas: 0.065 ml of He/min

No starting material was found.

5

Example 7:

Hydroxylation of aromatics, heteroaromatics and trimethylcyclohexenyl compounds

10

a) Example 6 was repeated, but using, instead of n-octane, the substrate naphthalene. The products that were identified were 1-naphthol and cis-1,2-dihydroxy-1,2-dihydronaphthalene. 88% of the naphthalene starting material had been converted.

15

Analytic methods for reactions with naphthalene

GC:

20

Apparatus: Carlo Erba Strumentazion Typ HRGC 4160 on Column
Injector; Column: DB5 30m x 0.2 mm; Material: 5% diphenyl-
95% dimethylpolysiloxane; Carrier gas: 0.5 bar H₂;
Temperature program: 40°C 1 min isothermic / 10°C/min to 300°C
Rt(1-naphthol) = 16.68

25

NMR:

1-Naphthol and cis-1,2-dihydroxy-1,2-dihydro-naphthalene were identified in the ¹H NMR.

30

b) Example 6 was repeated but using, instead of n-octane, the substrate 8-methylquinoline. 5-Hydroxy-8-methylquinoline was identified as main product, in addition to other derivatives (product ratio 5:1). 35% of the starting material used had been converted.

35

c) Example 6 was repeated but using, instead of n-octane, the substrate α -ionone. 3-Hydroxy- α -ionone was identified as main product, in addition to other derivatives (product ratio: 76:24). 60% of the starting material used had been converted.

40

d) Example 6 was repeated, but using, instead of n-octane, the substrate cumene (isopropylbenzene). Five monohydroxy products and one dihydroxy product were identified. 70% of the starting material used had been converted.

45

23

REFERENCES

1. Yano, T., Oue, S., and Kagamiyama, H. (1998) Proc. Natl. Acad Sci. USA 95, 5511-5515.
- 5 2. Zhang, J.-H., Dawes, G., and Stemmer, W. P. C. (1997) Proc. Natl. Acad Sci. USA 94, 4504-4509.
3. Wan, L., Twitchett, M. B., Eltis, L. D., Mauk, A. G., and
10 Smith, M. (1998) Proc. Natl. Acad Sci USA 95, 12825-12831.
4. Cronin, C. N. (1998) J. Biol. Chem 273, 24465-24469.
5. Wilks, H. M., Hart, K. W., Feeney, R., Dunn, C. R., Muirhead,
15 H., Chia, W. N., Barstow, D. A., Atkinson, T., Clarke, A. R.,
Holbrook, I J. (1988) Science 242, 1541-1544.
6. Hedstrom, L., Szilagyi, L., Rutter, W. J. (1992) Science 255,
1249-1253.
- 20 7. Tucker, C. L., Hurley, J. H., Miller, T. R., and Hurley, I B.
(1998) Proc. Natl. Acad Sci. USA 95, 5993-5997.
8. Quemeneur, E., Moutiez, J.-B. C., and Menez, A. (1998) Nature
25 (London) 391, 301-303.
9. Marsden, A- F. A., Wilkinson, B., Cortes, J., Dunster, N. J.,
Staunton, I Leadlay, P. F. (1998) Science 279, 199-201.
- 30 10. Chen, R., Greer, A., and Dean, A. M. (1998) Proc. Natl. Acad
Sci. US4 95, 11666-11670.
11. Boddupalli, S. S., Estabrook, R. W. and Peterson, J. A.
(1990) J Biol. Chem 265, 4233-4239.
- 35 12. Capdevila, J. H., Wie, S., Helvig, C., Falck, J. R.,
Belosludtsev, Y., Truan, G., Graham-Lorence, S. E., and
Peterson, J. A. (1996) J. Biol. Chem 271, 22663-22671.
- 40 13. Graham-Lorence, S., Truan, G., Peterson, J. A., Flack, J. R.,
WeL S., Helvig, C., Capdevilla, J. H. (1997) J. Biol. Chem
272, 1127-1135.
14. Li, H., Poulos, T. L. (1997) Nat. Structural Biol., 4,
45 140-146.

0050/50915

24

15. Ravichandran, K G., Sekhar, S., Boddupalli, S., Hasemann, C. A., Peterson, J. A., Deisenhofer, I (1993) Science 261, 731-736.
- 5 16. Modi S., Sutcliffe, M. J., Primrose, W. U., Lian, L.- Y., Roberts, G. C. K (1996) Nat. Structural Biol. 3, 414-417.
17. Oliver, C.F., Modi S., Primrose, W.U., Lian, L.Y. and Roberts, G.C.K (1997) Biochem. J. 327, 537-544.
- 10 18. Guengerich, F.G. (1991) J. Biol. Chem 266,10019-10022.
19. Cherry, J. R., Lamsa, M. H., Schneider, P., Vind, J., Svendsen, A-, Jones, A., and Pedersen, A. H. (1999) Nature Biotechnology 17, 379-384.
- 15 20. Schwaneberg, U., Schmidt-Dannert, C., Schmitt, J., and Schmid, R. D. (1999) Anal. Biochem. 269, 359-366.
- 20 21. Schwaneberg, U, Sprauer, AL, Schmidt-Dannert, C., and Schmid, R. D. J of Chromatogr. A, in press.
22. Oliver, C.F., Modi, S., Sutcliffe, M.J., Primrose, W.U., Lian, L.Y. and Roberts, G.C.K (1997) Biochemistry 36, 1567-1572.
- 25 23. Hart, S., Koch, KR., and Woods, D.R. (1992) J Gen. Microbiol. 138, 211-216
- 30 24. Murdock, D., Ensley, B.D., Serdar, C. and Thalen, M. (1993) Bio/Technology 11, 381-385.
25. O'Connor, ICE., Dobson, A-W. and Hartmans, S. (1997) Appl. Environ. Microbiol. 63, 4287-4291.
- 35 26. Eaton, R. W. and Chapman, P. J. (1995) J Bacteriol. 177, 6983-6988.

40

45

10/031146

JC13 Rec'd PCT/PTO 17 JAN 2002

1

We claim:

1. A cytochrome P450 monooxygenase which is capable of at least
5 one of the following reactions:
 - a) oxidation of optionally substituted N-, O- or
S-heterocyclic mono- or polynuclear aromatic compounds;
 - b) oxidation of optionally substituted mono- or polynuclear
10 aromatics;
 - c) oxidation of straight-chain or branched alkanes and
alkenes;
 - d) oxidation of optionally substituted cycloalkanes and
cycloalkenes;
- 15 where the monooxygenase is derived from cytochrome P450
monooxygenase BM-3 from *Bacillus megaterium* having an amino
acid sequence according to SEQ ID NO:2, which has at least
one functional mutation in at least one of the amino acid
20 sequence regions 172-224, 39-43, 48-52, 67-70, 330-335,
352-356, 73-82 and 86-88; except the single mutant Phe87Val.
2. A monooxygenase as claimed in claim 1, which has at least one
functional mutation in at least one of the sequence regions
73-82, 86-88 and 172-224.
- 25 3. A monooxygenase as claimed in claim 1, which has at least one
of the following mono- or polyamino acid substitutions:
 - a) Phe87Val, Leu188Gln; or
 - b) Phe87Val, Leu188Gln, Ala74Gly;
- 30 and functional equivalents thereof which are capable of at
least one of the above oxidation reactions.
4. A nucleic acid sequence coding for a monooxygenase according
to one of the preceding claims.
- 35 5. An expression construct comprising, under the genetic control
of regulatory nucleic acid sequences, a coding sequence which
comprises a nucleic acid sequence according to claim 4.
- 40 6. A vector comprising at least one expression construct
according to claim 5.
7. A recombinant microorganism transformed by at least one
45 vector as claimed in claim 6.

AMENDED SHEET

0050/50915

2

8. A microorganism as claimed in claim 7, selected from bacteria of the genus *Escherichia*.
9. A process for the microbiological oxidation of an N- or S-heterocyclic mono- or polynuclear aromatic compound, which comprises
- 5 a1) culturing a recombinant microorganism which expresses a cytochrome P450 monooxygenase of bacterial origin in a culture medium, in the presence of an exogenous or
- 10 a2) incubating a substrate-containing reaction medium with a cytochrome P450 monooxygenase of bacterial origin; and
- b) isolating the oxidation product formed or a secondary product thereof from the medium.
- 15 10. A process as claimed in claim 9, wherein the exogenous or intermediately formed substrate is selected from optionally substituted N- or S-heterocyclic mono- or polynuclear aromatic compounds.
- 20 11. A process as claimed in claim 9 or 10, where the monooxygenase is a mutant as claimed in any of claims 1 to 3, including the mutant Phe87Val.
- 25 12. A process as claimed in claim 11, where the mutant has at least one of the following mono- or polyamino acid substitutions:
- a) Phe87Val;
- b) Phe87Val, Leu188Gln; or
- 30 c) Phe87Val, Leu188Gln, Ala74Gly.
13. A process for microbiological oxidation of a compound as defined in claim 1b), c) or d), which comprises
- 35 a1) culturing a recombinant cytochrome P450-producing microorganism as claimed in claim 7 or 8 in a culture medium, in the presence of an exogenous or intermediately formed substrate; or
- 40 a2) incubating a substrate-containing reaction medium with a cytochrome P450 monooxygenase as claimed in any of claims 1 to 3; and
- b) isolating the oxidation product formed or a secondary product thereof from the medium;
- 45 where the monooxygenase mutant Phe87Val is not excluded.
14. A process as claimed in claim 13, wherein the exogenous or intermediately formed substrate is selected from:

AMENDED SHEET

3

- a) optionally substituted mono- or polynuclear aromatics;
- b) straight-chain or branched alkanes and alkenes;
- c) optionally substituted cycloalkanes and cycloalkenes.

5

15. A process as claimed in claim 13 or 14, where the monooxygenase is a mutant as claimed in any of claims 1 to 3, including the mutant Phe87Val.

10 16. A process as claimed in claim 15, where the mutant has at least one of the following mono- or polyamino acid substitutions:

- a) Phe87Val;
- b) Phe87Val, Leu188Gln; or
- 15 c) Phe87Val, Leu188Gln, Ala74Gly.

17. A process as claimed in any of claims 9 to 16, wherein, as exogenous substrate, at least one compound selected from the groups a) to d) of compounds defined above is added to a
20 medium and the oxidation is carried out by enzymatic reaction of the substrate-containing medium in the presence of oxygen at a temperature of approximately 20 to 40°C and a pH of approximately 6 to 9, where the substrate-containing medium additionally contains an approximately 10- to 100-fold molar
25 excess of reduction equivalents based on the substrate.

18. A process as claimed in claim 17, wherein, as exogenous substrate, a compound selected from indole, n-hexane, n-octane, n-decane, n-dodecane, cumene, 1-methylindole, α -, β -
30 or γ -ionone, acridine, naphthalene, 6-methyl- or 8-methylquinoline, quinoline and quinaldine is employed.

19. A process for the microbiological production of indigo and/or indirubin, which comprises
35

- a1) culturing a recombinant microorganism which produces an indole-oxidizing cytochrome P450 in a culture medium, in the presence of exogenous or intermediately formed indole; or
- a2) incubating an indole-containing reaction medium with an
40 indole-oxidizing cytochrome P450 monooxygenase; and
- b) isolating the oxidation product formed or a secondary product thereof from the medium;

20. A process as claimed in claim 19, wherein the indigo and/or indirubin obtained, which was produced by oxidation of
45 intermediately formed indole, is isolated from the medium.

AMENDED SHEET

4

- 45

AMENDED SHEET

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG(19) Weltorganisation für geistiges Eigentum
Internationales Büro(43) Internationales Veröffentlichungsdatum
1. Februar 2001 (01.02.2001)

PCT

(10) Internationale Veröffentlichungsnummer
WO 01/07630 A1(51) Internationale Patentklassifikation⁷: **C12N 15/53**,
9/02, 15/70, 1/21, C12P 17/10, 17/16, 7/04, 7/22, 7/02
// (C12N 1/21, C12R 1:19)**Qing-shan** [JP/JP]; Kitashirakawa-oiwakecho, Sakyo-ku,
Kyoto 606-8502 (JP).

(21) Internationales Aktenzeichen: PCT/EP00/07253

(74) Anwälte: **KINZEBACH, Werner** usw.; Reitstötter,
Kinzebach & Partner, Sternwartstrasse 4, D-81679
München (DE).(22) Internationales Anmeldedatum:
27. Juli 2000 (27.07.2000)(81) Bestimmungsstaaten (*national*): AE, AG, AL, AM, AT,
AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO,
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:
199 35 115.5 27. Juli 1999 (27.07.1999) DE
199 55 605.9 18. November 1999 (18.11.1999) DE
100 14 085.8 22. März 2000 (22.03.2000) DE(84) Bestimmungsstaaten (*regional*): ARIPO-Patent (GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), eura-
sisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI,
FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI-Patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE,
SN, TD, TG).(71) Anmelder (*für alle Bestimmungsstaaten mit Ausnahme
von US*): **BASF AKTIENGESSELLSCHAFT** [DE/DE];
D-67056 Ludwigshafen (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder (*nur für US*): **HAUER, Bernhard**
[DE/DE]; Merowingerstrasse 1, D-67136 Fussgönheim
(DE). **PLEISS, Juergen** [DE/DE]; Ostlandstrasse 14,
D-71679 Asperg (DE). **SCHWANEGER, Ulrich**
[DE/DE]; Uhlandstrasse 15, D-71336 Waiblingen (DE).
SCHMITT, Jutta [DE/DE]; Fuggerstrasse 19, D-70563
Stuttgart (DE). **FISCHER, Markus** [DE/DE]; Uhland-
strasse 14, D-71638 Ludwigsburg (DE). **SCHMID, Rolf**
[DE/DE]; Sylvanerweg 6, D-79329 Stuttgart (DE). **LI,**

Veröffentlicht:

- Mit internationalem Recherchenbericht.
- Vor Ablauf der für Änderungen der Ansprüche geltenden Frist; Veröffentlichung wird wiederholt, falls Änderungen eintreffen.

Zur Erklärung der Zweibuchstaben-Codes, und der anderen
Abkürzungen wird auf die Erklärungen ("Guidance Notes on
Codes and Abbreviations") am Anfang jeder regulären Ausgabe
der PCT-Gazette verwiesen.(54) Title: NOVEL CYTOCHROME P450 MONOOXYGENASES AND THEIR USE FOR OXIDIZING ORGANIC
COMPOUNDS(54) Bezeichnung: NEUE CYTOCHROM P450-MONOOXYGENASEN UND DEREN VERWENDUNG ZUR OXIDATION
VON ORGANISCHEN VERBINDUNGEN(57) Abstract: The invention relates to novel cytochrome P450 monooxygenases comprising a modified substrate specificity, to
nucleotide sequences which code therefor, to expression constructs and vectors containing these sequences, and to microorganisms
transformed therewith. The invention also relates to methods for microbiologically oxidizing different organic substrates, such as
methods for producing indigo and indirubin.(57) Zusammenfassung: Die Erfindung betrifft neue Cytochrom P450-Monooxygenasen mit veränderter Substratspezifität, dafür
kodierende Nukleotidsequenzen, diese Sequenzen enthaltende Expressionskonstrukte und Vektoren, damit transformierte Mikroor-
ganismen, Verfahren zur mikrobiologischen Oxidation verschiedener organischer Substrate wie beispielsweise Verfahren zur Her-
stellung von Indigo und Indirubin.

WO 01/07630 A1

Declaration, Power of Attorney and Petition

Page 1 of 4
0050/050915

We (I), the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Novel cytochrome P450 monooxygenases and their use for oxidizing organic compounds

the specification of which

☒ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and amended on _____.

☒ was filed as PCT international application

Number PCT/EP/00/07253

on 27 July 2000,

and was amended under PCT Article 19

on _____ (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)–(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed
19935115.5	Germany	27 July 1999	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
19955605.9	Germany	18 November 1999	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
10014085.5	Germany	22 March 2000	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No

Declaration

Page 2 of 4

0050/050915

We (I) hereby claim the benefit under Title 35, United States Codes, § 119(e) of any United States provisional application(s) listed below.

_____	_____
(Application Number)	(Filing Date)
_____	_____
(Application Number)	(Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.	Filing Date	Status (pending, patented, abandoned)
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

(2)

And we (I) hereby appoint Messrs. **HERBERT B. KEIL**, Registration Number 18,967; and **RUSSEL E. WEINKAUF**, Registration Number 18,495; the address of both being Messrs. Keil & Weinkauff, 1101 Connecticut Ave., N.W., Washington, D.C. 20036 (telephone 202-659-0100), our attorneys, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to sign the drawings, to receive the patent, and to transact all business in the Patent Office connected therewith.

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Declaration

Page 3 of 4

0050/050915

1-00
Bernhard Hauer
NAME OF INVENTOR

Bernhard Hauer
Signature of Inventor

Date 27 July 2000

Merowingerstr. 1
67056 Fußgönheim DEX
Germany
Citizen of: Germany
Post Office Address: same as residence

2-00
Juergen Pleiss
NAME OF INVENTOR

Juergen Pleiss
Signature of Inventor

Date 27 July 2000

Ostlandstr. 13 DEX
71679 Asperg
Germany
Citizen of: Germany
Post Office Address: same as residence

3-00
Ulrich Schwaneberg
NAME OF INVENTOR

Ulrich Schwaneberg
Signature of Inventor

Date 27 July 2000

Uhlandstr. 15 DEX
71336 Waiblingen
Germany
Citizen of: Germany
Post Office Address: same as residence

4-00
Jutta Schmitt
NAME OF INVENTOR

Jutta Schmitt
Signature of Inventor

Date 27 July 2000

Fuggerstr. 19 DEX
70563 Stuttgart
Germany
Citizen of: Germany
Post Office Address: same as residence

Declaration

Page 4 of 4

0050/050915

5-00
Markus Fischer
 NAME OF INVENTOR

Markus Fischer
 Signature of Inventor

Date 27 July 2000

Uhlandstr. 14
 71638 Ludwigsburg DEX
 Germany
 Citizen of: Germany
 Post Office Address: same as residence

6-00
Rolf Schmid
 NAME OF INVENTOR

Rolf Schmid
 Signature of Inventor

Date 27 July 2000

Sylvanerweg 6
 70329 Stuttgart DEX
 Germany
 Citizen of: Germany
 Post Office Address: same as residence

7-00
Qing-shan Li
 NAME OF INVENTOR

Qingshan Li
 Signature of Inventor

Date 27 July 2000

Kitashirakawa-oiwakecho
 Sakyo-ku, Kyoto 606-8502
 Japan Qingshan Li CEX
 Citizen of: ~~Japan~~ China
 Post Office Address: same as residence

8-00
Sabine Lutz-Wahl
 NAME OF INVENTOR

Sabine Lutz-Wahl
 Signature of Inventor

Date: 27 July 2000

Bogenstr. 41
 70569 Stuttgart DEX
 Germany
 Citizen of: Germany
 Post Office Address: same as residence

9-00
Daniel Appel
 NAME OF INVENTOR

Daniel Appel
 SIGNATURE of INVENTOR

Date: 27 July 2000

August-Lämmle-Weg 5
 74348 Lauffen DEX
 Germany
 Citizen of: Germany
 Post Office Address: same as residence

10/031146
PCT Rec'd 30 APR 2002

1

SEQUENCE LISTING

<110> HAUER, Bernhard
 PLEISS, Juergen
 SCHWANEBERG, Ulrich
 SCHMITT, Jutta

<120> Novel cytochrome P450 monooxygenases and their use for the oxidation of organic substrates

<130> 50915

<140> 10/031,146

<141> Filing date not yet granted

<150> PCT/EP 00/07253

<151> 2000-07-27

<160> 9

<170> Word Perfect version 6.1

<210> 1

<211> 3150

<212> DNA

<213> Bacillus megaterium

<220>

<221> CDS

<222> (4)..(3150)

<400> 1

atg aca att aaa gaa atg cct cag cca aaa acg ttt gga gag ctt aaa	48
Thr Ile Lys Glu Met Pro Gln Pro Lys Thr Phe Gly Glu Leu Lys	
1 5 10 15	
aat tta ccg tta tta aac aca gat aaa ccg gtt caa gct ttg atg aaa	96
Asn Leu Pro Leu Leu Asn Thr Asp Lys Pro Val Gln Ala Leu Met Lys	
20 25 30	
att gcg gat gaa tta gga gaa atc ttt aaa ttc gag gcg cct ggt cgt	144
Ile Ala Asp Glu Leu Gly Glu Ile Phe Lys Phe Glu Ala Pro Gly Arg	
35 40 45	
gta acg cgc tac tta tca agt cag cgt cta att aaa gaa gca tgc gat	192
Val Thr Arg Tyr Leu Ser Ser Gln Arg Leu Ile Lys Glu Ala Cys Asp	
50 55 60	
gaa tca cgc ttt gat aaa aac tta agt caa gcg ctt aaa ttt gta cgt	240
Glu Ser Arg Phe Asp Lys Asn Leu Ser Gln Ala Leu Lys Phe Val Arg	
65 70 75	
gat ttt gca gga gac ggg tta ttt aca agc tgg acg cat gaa aaa aat	288
Asp Phe Ala Gly Asp Gly Leu Phe Thr Ser Trp Thr His Glu Lys Asn	
80 85 90 95	
tgg aaa aaa gcg cat aat atc tta ctt cca agc ttc agt cag cag gca	336
Trp Lys Lys Ala His Asn Ile Leu Leu Pro Ser Phe Ser Gln Gln Ala	

	100	105	110	
atg aaa ggc tat cat gcg atg atg gtc gat atc gcc gtg cag ctt gtt	384			
Met Lys Gly Tyr His Ala Met Met Val Asp Ile Ala Val Gln Leu Val				
	115	120	125	
caa aag tgg gag cgt cta aat gca gat gag cat att gaa gta ccg gaa	432			
Gln Lys Trp Glu Arg Leu Asn Ala Asp Glu His Ile Glu Val Pro Glu				
	130	135	140	
gac atg aca cgt tta acg ctt gat aca att ggt ctt tgc ggc ttt aac	480			
Asp Met Thr Arg Leu Thr Leu Asp Thr Ile Gly Leu Cys Gly Phe Asn				
	145	150	155	
tat cgc ttt aac agc ttt tac cga gat cag cct cat cca ttt att aca	528			
Tyr Arg Phe Asn Ser Phe Tyr Arg Asp Gln Pro His Pro Phe Ile Thr				
	160	165	170	175
agt atg gtc cgt gca ctg gat gaa gca atg aac aag ctg cag cga gca	576			
Ser Met Val Arg Ala Leu Asp Glu Ala Met Asn Lys Leu Gln Arg Ala				
	180	185	190	
aat cca gac gac cca gct tat gat gaa aac aag cgc cag ttt caa gaa	624			
Asn Pro Asp Asp Pro Ala Tyr Asp Glu Asn Lys Arg Gln Phe Gln Glu				
	195	200	205	
gat atc aag gtg atg aac gac cta gta gat aaa att att gca gat cgc	672			
Asp Ile Lys Val Met Asn Asp Leu Val Asp Lys Ile Ile Ala Asp Arg				
	210	215	220	
aaa gca agc ggt gaa caa agc gat gat tta tta acg cat atg cta aac	720			
Lys Ala Ser Gly Glu Gln Ser Asp Asp Leu Leu Thr His Met Leu Asn				
	225	230	235	
gga aaa gat cca gaa acg ggt gag ccg ctt gat gac gag aac att cgc	768			
Gly Lys Asp Pro Glu Thr Gly Glu Pro Leu Asp Asp Glu Asn Ile Arg				
	240	245	250	255
tat caa att att aca ttc tta att gcg gga cac gaa aca aca agt ggt	816			
Tyr Gln Ile Ile Thr Phe Leu Ile Ala Gly His Glu Thr Thr Ser Gly				
	260	265	270	
ctt tta tca ttt gcg ctg tat ttc tta gtg aaa aat cca cat gta tta	864			
Leu Leu Ser Phe Ala Leu Tyr Phe Leu Val Lys Asn Pro His Val Leu				
	275	280	285	
caa aaa gca gca gaa gaa gca gca cga gtt cta gta gat cct gtt cca	912			
Gln Lys Ala Ala Glu Glu Ala Ala Arg Val Leu Val Asp Pro Val Pro				
	290	295	300	
agc tac aaa caa gtc aaa cag ctt aaa tat gtc ggc atg gtc tta aac	960			
Ser Tyr Lys Gln Val Lys Gln Leu Lys Tyr Val Gly Met Val Leu Asn				
	305	310	315	
gaa gcg ctg cgc tta tgg cca act gct cct gcg ttt tcc cta tat gca	1008			
Glu Ala Leu Arg Leu Trp Pro Thr Ala Pro Ala Phe Ser Leu Tyr Ala				
	320	325	330	335
aaa gaa gat acg gtg ctt gga gga gaa tat cct tta gaa aaa ggc gac	1056			

Lys	Glu	Asp	Thr	Val	Leu	Gly	Gly	Glu	Tyr	Pro	Leu	Glu	Lys	Gly	Asp		
				340					345					350			
gaa	cta	atg	gtt	ctg	att	cct	cag	ctt	cac	cgt	gat	aaa	aca	att	tgg	1104	
Glu	Leu	Met	Val	Leu	Ile	Pro	Gln	Leu	His	Arg	Asp	Lys	Thr	Ile	Trp		
			355					360					365				
gga	gac	gat	gtg	gaa	gag	ttc	cgt	cca	gag	cgt	ttt	gaa	aat	cca	agt	1152	
Gly	Asp	Asp	Val	Glu	Glu	Phe	Arg	Pro	Glu	Arg	Phe	Glu	Asn	Pro	Ser		
			370				375					380					
gcg	att	ccg	cag	cat	gcg	ttt	aaa	ccg	ttt	gga	aac	ggg	cag	cgt	gcg	1200	
Ala	Ile	Pro	Gln	His	Ala	Phe	Lys	Pro	Phe	Gly	Asn	Gly	Gln	Arg	Ala		
	385					390				395							
tgt	atc	ggg	cag	cag	ttc	gct	ctt	cat	gaa	gca	acg	ctg	gta	ctt	ggg	1248	
Cys	Ile	Gly	Gln	Gln	Phe	Ala	Leu	His	Glu	Ala	Thr	Leu	Val	Leu	Gly		
400				405					410						415		
atg	atg	cta	aaa	cac	ttt	gac	ttt	gaa	gat	cat	aca	aac	tac	gag	ctg	1296	
Met	Met	Leu	Lys	His	Phe	Asp	Phe	Glu	Asp	His	Thr	Asn	Tyr	Glu	Leu		
				420				425						430			
gat	att	aaa	gaa	act	tta	acg	tta	aaa	cct	gaa	ggc	ttt	gtg	gta	aaa	1344	
Asp	Ile	Lys	Glu	Thr	Leu	Thr	Leu	Lys	Pro	Glu	Gly	Phe	Val	Val	Lys		
			435				440						445				
gca	aaa	tgc	aaa	aaa	att	ccg	ctt	ggc	ggg	att	cct	tca	cct	agc	act	1392	
Ala	Lys	Ser	Lys	Lys	Ile	Pro	Leu	Gly	Gly	Ile	Pro	Ser	Pro	Ser	Thr		
		450					455					460					
gaa	cag	tct	gct	aaa	aaa	gta	cgc	aaa	aag	gca	gaa	aac	gct	cat	aat	1440	
Glu	Gln	Ser	Ala	Lys	Lys	Val	Arg	Lys	Lys	Ala	Glu	Asn	Ala	His	Asn		
	465					470				475							
acg	ccg	ctg	ctt	gtg	cta	tac	ggg	tca	aat	atg	gga	aca	gct	gaa	gga	1488	
Thr	Pro	Leu	Leu	Val	Leu	Tyr	Gly	Ser	Asn	Met	Gly	Thr	Ala	Glu	Gly		
480				485				490							495		
acg	gcg	cgt	gat	tta	gca	gat	att	gca	atg	agc	aaa	gga	ttt	gca	ccg	1536	
Thr	Ala	Arg	Asp	Leu	Ala	Asp	Ile	Ala	Met	Ser	Lys	Gly	Phe	Ala	Pro		
				500				505						510			
cag	gtc	gca	acg	ctt	gat	tca	cac	gcc	gga	aat	ctt	ccg	cgc	gaa	gga	1584	
Gln	Val	Ala	Thr	Leu	Asp	Ser	His	Ala	Gly	Asn	Leu	Pro	Arg	Glu	Gly		
			515					520					525				
gct	gta	tta	att	gta	acg	gcg	tct	tat	aac	ggg	cat	ccg	cct	gat	aac	1632	
Ala	Val	Leu	Ile	Val	Thr	Ala	Ser	Tyr	Asn	Gly	His	Pro	Pro	Asp	Asn		
		530					535					540					
gca	aag	caa	ttt	gtc	gac	tgg	tta	gac	caa	gcg	tct	gct	gat	gaa	gta	1680	
Ala	Lys	Gln	Phe	Val	Asp	Trp	Leu	Asp	Gln	Ala	Ser	Ala	Asp	Glu	Val		
	545					550				555							
aaa	ggc	gtt	cgc	tac	tcc	gta	ttt	gga	tgc	ggc	gat	aaa	aac	tgg	gct	1728	
Lys	Gly	Val	Arg	Tyr	Ser	Val	Phe	Gly	Cys	Gly	Asp	Lys	Asn	Trp	Ala		
560				565				570							575		

act	acg	tat	caa	aaa	gtg	cct	gct	ttt	atc	gat	gaa	acg	ctt	gcc	gct	1776
Thr	Thr	Tyr	Gln	Lys	Val	Pro	Ala	Phe	Ile	Asp	Glu	Thr	Leu	Ala	Ala	
				580					585					590		
aaa	ggg	gca	gaa	aac	atc	gct	gac	cgc	ggg	gaa	gca	gat	gca	agc	gac	1824
Lys	Gly	Ala	Glu	Asn	Ile	Ala	Asp	Arg	Gly	Glu	Ala	Asp	Ala	Ser	Asp	
			595					600					605			
gac	ttt	gaa	ggc	aca	tat	gaa	gaa	tgg	cgt	gaa	cat	atg	tgg	agt	gac	1872
Asp	Phe	Glu	Gly	Thr	Tyr	Glu	Glu	Trp	Arg	Glu	His	Met	Trp	Ser	Asp	
		610						615				620				
gta	gca	gcc	tac	ttt	aac	ctc	gac	att	gaa	aac	agt	gaa	gat	aat	aaa	1920
Val	Ala	Ala	Tyr	Phe	Asn	Leu	Asp	Ile	Glu	Asn	Ser	Glu	Asp	Asn	Lys	
	625					630					635					
tct	act	ctt	tca	ctt	caa	ttt	gtc	gac	agc	gcc	gcg	gat	atg	ccg	ctt	1968
Ser	Thr	Leu	Ser	Leu	Gln	Phe	Val	Asp	Ser	Ala	Ala	Asp	Met	Pro	Leu	
640					645					650					655	
gcg	aaa	atg	cac	ggg	gcg	ttt	tca	acg	aac	gtc	gta	gca	agc	aaa	gaa	2016
Ala	Lys	Met	His	Gly	Ala	Phe	Ser	Thr	Asn	Val	Val	Ala	Ser	Lys	Glu	
				660					665					670		
ctt	caa	cag	cca	ggc	agt	gca	cga	agc	acg	cga	cat	ctt	gaa	att	gaa	2064
Leu	Gln	Gln	Pro	Gly	Ser	Ala	Arg	Ser	Thr	Arg	His	Leu	Glu	Ile	Glu	
			675					680					685			
ctt	cca	aaa	gaa	gct	tct	tat	caa	gaa	gga	gat	cat	tta	ggg	gtt	att	2112
Leu	Pro	Lys	Glu	Ala	Ser	Tyr	Gln	Glu	Gly	Asp	His	Leu	Gly	Val	Ile	
		690					695					700				
cct	cgc	aac	tat	gaa	gga	ata	gta	aac	cgt	gta	aca	gca	agg	ttc	ggc	2160
Pro	Arg	Asn	Tyr	Glu	Gly	Ile	Val	Asn	Arg	Val	Thr	Ala	Arg	Phe	Gly	
	705					710					715					
cta	gat	gca	tca	cag	caa	atc	cgt	ctg	gaa	gca	gaa	gaa	gaa	aaa	tta	2208
Leu	Asp	Ala	Ser	Gln	Gln	Ile	Arg	Leu	Glu	Ala	Glu	Glu	Glu	Lys	Leu	
720					725					730					735	
gct	cat	ttg	cca	ctc	gct	aaa	aca	gta	tcc	gta	gaa	gag	ctt	ctg	caa	2256
Ala	His	Leu	Pro	Leu	Ala	Lys	Thr	Val	Ser	Val	Glu	Glu	Leu	Leu	Gln	
				740					745					750		
tac	gtg	gag	ctt	caa	gat	cct	gtt	acg	cgc	acg	cag	ctt	cgc	gca	atg	2304
Tyr	Val	Glu	Leu	Gln	Asp	Pro	Val	Thr	Arg	Thr	Gln	Leu	Arg	Ala	Met	
			755					760					765			
gct	gct	aaa	acg	gtc	tgc	ccg	ccg	cat	aaa	gta	gag	ctt	gaa	gcc	ttg	2352
Ala	Ala	Lys	Thr	Val	Cys	Pro	Pro	His	Lys	Val	Glu	Leu	Glu	Ala	Leu	
		770					775					780				
ctt	gaa	aag	caa	gcc	tac	aaa	gaa	caa	gtg	ctg	gca	aaa	cgt	tta	aca	2400
Leu	Glu	Lys	Gln	Ala	Tyr	Lys	Glu	Gln	Val	Leu	Ala	Lys	Arg	Leu	Thr	
	785					790					795					
atg	ctt	gaa	ctg	ctt	gaa	aaa	tac	ccg	gcg	tgt	gaa	atg	aaa	ttc	agc	2448
Met	Leu	Glu	Leu	Leu	Glu	Lys	Tyr	Pro	Ala	Cys	Glu	Met	Lys	Phe	Ser	
800					805					810					815	

gaa ttt atc gcc ctt ctg cca agc ata cgc ccg cgc tat tac tcg att	2496
Glu Phe Ile Ala Leu Pro Ser Ile Arg Pro Arg Tyr Tyr Ser Ile	
820 825 830	
tct tca tca cct cgt gtc gat gaa aaa caa gca agc atc acg gtc agc	2544
Ser Ser Ser Pro Arg Val Asp Glu Lys Gln Ala Ser Ile Thr Val Ser	
835 840 845	
gtt gtc tca gga gaa gcg tgg agc gga tat gga gaa tat aaa gga att	2592
Val Val Ser Gly Glu Ala Trp Ser Gly Tyr Gly Glu Tyr Lys Gly Ile	
850 855 860	
gcg tcg aac tat ctt gcc gag ctg caa gaa gga gat acg att acg tgc	2640
Ala Ser Asn Tyr Leu Ala Glu Leu Gln Glu Gly Asp Thr Ile Thr Cys	
865 870 875	
ttt att tcc aca ccg cag tca gaa ttt acg ctg cca aaa gac cct gaa	2688
Phe Ile Ser Thr Pro Gln Ser Glu Phe Thr Leu Pro Lys Asp Pro Glu	
880 885 890 895	
acg ccg ctt atc atg gtc gga ccg gga aca ggc gtc gcg ccg ttt aga	2736
Thr Pro Leu Ile Met Val Gly Pro Gly Thr Gly Val Ala Pro Phe Arg	
900 905 910	
ggc ttt gtg cag gcg cgc aaa cag cta aaa gaa caa gga cag tca ctt	2784
Gly Phe Val Gln Ala Arg Lys Gln Leu Lys Glu Gln Gly Gln Ser Leu	
915 920 925	
gga gaa gca cat tta tac ttc ggc tgc cgt tca cct cat gaa gac tat	2832
Gly Glu Ala His Leu Tyr Phe Gly Cys Arg Ser Pro His Glu Asp Tyr	
930 935 940	
ctg tat caa gaa gag ctt gaa aac gcc caa agc gaa ggc atc att acg	2880
Leu Tyr Gln Glu Glu Leu Glu Asn Ala Gln Ser Glu Gly Ile Ile Thr	
945 950 955	
ctt cat acc gct ttt tct cgc atg cca aat cag ccg aaa aca tac gtt	2928
Leu His Thr Ala Phe Ser Arg Met Pro Asn Gln Pro Lys Thr Tyr Val	
960 965 970 975	
cag cac gta atg gaa caa gac ggc aag aaa ttg att gaa ctt ctt gat	2976
Gln His Val Met Glu Gln Asp Gly Lys Lys Leu Ile Glu Leu Leu Asp	
980 985 990	
caa gga gcg cac ttc tat att tgc gga gac gga agc caa atg gca cct	3024
Gln Gly Ala His Phe Tyr Ile Cys Gly Asp Gly Ser Gln Met Ala Pro	
995 1000 1005	
gcc gtt gaa gca acg ctt atg aaa agc tat gct gac gtt cac caa gtg	3072
Ala Val Glu Ala Thr Leu Met Lys Ser Tyr Ala Asp Val His Gln Val	
1010 1015 1020	
agt gaa gca gac gct cgc tta tgg ctg cag cag cta gaa gaa aaa ggc	3120
Ser Glu Ala Asp Ala Arg Leu Trp Leu Gln Gln Leu Glu Glu Lys Gly	
1025 1030 1035	
cga tac gca aaa gac gtg tgg gct ggg taa	3150
Arg Tyr Ala Lys Asp Val Trp Ala Gly	
1040 1045	

<210> 2
 <211> 1048
 <212> PRT
 <213> Bacillus megaterium

<400> 2

Thr	Ile	Lys	Glu	Met	Pro	Gln	Pro	Lys	Thr	Phe	Gly	Glu	Leu	Lys	Asn	1	5	10	15
Leu	Pro	Leu	Leu	Asn	Thr	Asp	Lys	Pro	Val	Gln	Ala	Leu	Met	Lys	Ile	20	25	30	
Ala	Asp	Glu	Leu	Gly	Glu	Ile	Phe	Lys	Phe	Glu	Ala	Pro	Gly	Arg	Val	35	40	45	
Thr	Arg	Tyr	Leu	Ser	Ser	Gln	Arg	Leu	Ile	Lys	Glu	Ala	Cys	Asp	Glu	50	55	60	
Ser	Arg	Phe	Asp	Lys	Asn	Leu	Ser	Gln	Ala	Leu	Lys	Phe	Val	Arg	Asp	65	70	75	80
Phe	Ala	Gly	Asp	Gly	Leu	Phe	Thr	Ser	Trp	Thr	His	Glu	Lys	Asn	Trp	85	90	95	
Lys	Lys	Ala	His	Asn	Ile	Leu	Leu	Pro	Ser	Phe	Ser	Gln	Gln	Ala	Met	100	105	110	
Lys	Gly	Tyr	His	Ala	Met	Met	Val	Asp	Ile	Ala	Val	Gln	Leu	Val	Gln	115	120	125	
Lys	Trp	Glu	Arg	Leu	Asn	Ala	Asp	Glu	His	Ile	Glu	Val	Pro	Glu	Asp	130	135	140	
Met	Thr	Arg	Leu	Thr	Leu	Asp	Thr	Ile	Gly	Leu	Cys	Gly	Phe	Asn	Tyr	145	150	155	160
Arg	Phe	Asn	Ser	Phe	Tyr	Arg	Asp	Gln	Pro	His	Pro	Phe	Ile	Thr	Ser	165	170	175	
Met	Val	Arg	Ala	Leu	Asp	Glu	Ala	Met	Asn	Lys	Leu	Gln	Arg	Ala	Asn	180	185	190	
Pro	Asp	Asp	Pro	Ala	Tyr	Asp	Glu	Asn	Lys	Arg	Gln	Phe	Gln	Glu	Asp	195	200	205	
Ile	Lys	Val	Met	Asn	Asp	Leu	Val	Asp	Lys	Ile	Ile	Ala	Asp	Arg	Lys	210	215	220	
Ala	Ser	Gly	Glu	Gln	Ser	Asp	Asp	Leu	Leu	Thr	His	Met	Leu	Asn	Gly	225	230	235	240
Lys	Asp	Pro	Glu	Thr	Gly	Glu	Pro	Leu	Asp	Asp	Glu	Asn	Ile	Arg	Tyr	245	250	255	
Gln	Ile	Ile	Thr	Phe	Leu	Ile	Ala	Gly	His	Glu	Thr	Thr	Ser	Gly	Leu	260	265	270	

Leu Ser Phe Ala Leu Tyr Phe Leu Val Lys Asn Pro His Val Leu Gln
 275 280 285
 Lys Ala Ala Glu Glu Ala Ala Arg Val Leu Val Asp Pro Val Pro Ser
 290 295 300
 Tyr Lys Gln Val Lys Gln Leu Lys Tyr Val Gly Met Val Leu Asn Glu
 305 310 315 320
 Ala Leu Arg Leu Trp Pro Thr Ala Pro Ala Phe Ser Leu Tyr Ala Lys
 325 330 335
 Glu Asp Thr Val Leu Gly Gly Glu Tyr Pro Leu Glu Lys Gly Asp Glu
 340 345 350
 Leu Met Val Leu Ile Pro Gln Leu His Arg Asp Lys Thr Ile Trp Gly
 355 360 365
 Asp Asp Val Glu Glu Phe Arg Pro Glu Arg Phe Glu Asn Pro Ser Ala
 370 375 380
 Ile Pro Gln His Ala Phe Lys Pro Phe Gly Asn Gly Gln Arg Ala Cys
 385 390 395 400
 Ile Gly Gln Gln Phe Ala Leu His Glu Ala Thr Leu Val Leu Gly Met
 405 410 415
 Met Leu Lys His Phe Asp Phe Glu Asp His Thr Asn Tyr Glu Leu Asp
 420 425 430
 Ile Lys Glu Thr Leu Thr Leu Lys Pro Glu Gly Phe Val Val Lys Ala
 435 440 445
 Lys Ser Lys Lys Ile Pro Leu Gly Gly Ile Pro Ser Pro Ser Thr Glu
 450 455 460
 Gln Ser Ala Lys Lys Val Arg Lys Lys Ala Glu Asn Ala His Asn Thr
 465 470 475 480
 Pro Leu Leu Val Leu Tyr Gly Ser Asn Met Gly Thr Ala Glu Gly Thr
 485 490 495
 Ala Arg Asp Leu Ala Asp Ile Ala Met Ser Lys Gly Phe Ala Pro Gln
 500 505 510
 Val Ala Thr Leu Asp Ser His Ala Gly Asn Leu Pro Arg Glu Gly Ala
 515 520 525
 Val Leu Ile Val Thr Ala Ser Tyr Asn Gly His Pro Pro Asp Asn Ala
 530 535 540
 Lys Gln Phe Val Asp Trp Leu Asp Gln Ala Ser Ala Asp Glu Val Lys
 545 550 555 560
 Gly Val Arg Tyr Ser Val Phe Gly Cys Gly Asp Lys Asn Trp Ala Thr
 565 570 575
 Thr Tyr Gln Lys Val Pro Ala Phe Ile Asp Glu Thr Leu Ala Ala Lys
 580 585 590

Gly	Ala	Glu	Asn	Ile	Ala	Asp	Arg	Gly	Glu	Ala	Asp	Ala	Ser	Asp	Asp	
		595					600					605				
Phe	Glu	Gly	Thr	Tyr	Glu	Glu	Trp	Arg	Glu	His	Met	Trp	Ser	Asp	Val	
	610					615					620					
Ala	Ala	Tyr	Phe	Asn	Leu	Asp	Ile	Glu	Asn	Ser	Glu	Asp	Asn	Lys	Ser	
625					630					635					640	
Thr	Leu	Ser	Leu	Gln	Phe	Val	Asp	Ser	Ala	Ala	Asp	Met	Pro	Leu	Ala	
				645					650					655		
Lys	Met	His	Gly	Ala	Phe	Ser	Thr	Asn	Val	Val	Ala	Ser	Lys	Glu	Leu	
			660					665					670			
Gln	Gln	Pro	Gly	Ser	Ala	Arg	Ser	Thr	Arg	His	Leu	Glu	Ile	Glu	Leu	
		675					680					685				
Pro	Lys	Glu	Ala	Ser	Tyr	Gln	Glu	Gly	Asp	His	Leu	Gly	Val	Ile	Pro	
	690					695					700					
Arg	Asn	Tyr	Glu	Gly	Ile	Val	Asn	Arg	Val	Thr	Ala	Arg	Phe	Gly	Leu	
705					710					715					720	
Asp	Ala	Ser	Gln	Gln	Ile	Arg	Leu	Glu	Ala	Glu	Glu	Glu	Lys	Leu	Ala	
				725					730					735		
His	Leu	Pro	Leu	Ala	Lys	Thr	Val	Ser	Val	Glu	Glu	Leu	Leu	Gln	Tyr	
			740					745					750			
Val	Glu	Leu	Gln	Asp	Pro	Val	Thr	Arg	Thr	Gln	Leu	Arg	Ala	Met	Ala	
		755				760						765				
Ala	Lys	Thr	Val	Cys	Pro	Pro	His	Lys	Val	Glu	Leu	Glu	Ala	Leu	Leu	
	770					775					780					
Glu	Lys	Gln	Ala	Tyr	Lys	Glu	Gln	Val	Leu	Ala	Lys	Arg	Leu	Thr	Met	
785					790					795					800	
Leu	Glu	Leu	Leu	Glu	Lys	Tyr	Pro	Ala	Cys	Glu	Met	Lys	Phe	Ser	Glu	
				805					810					815		
Phe	Ile	Ala	Leu	Leu	Pro	Ser	Ile	Arg	Pro	Arg	Tyr	Tyr	Ser	Ile	Ser	
			820					825					830			
Ser	Ser	Pro	Arg	Val	Asp	Glu	Lys	Gln	Ala	Ser	Ile	Thr	Val	Ser	Val	
		835					840					845				
Val	Ser	Gly	Glu	Ala	Trp	Ser	Gly	Tyr	Gly	Glu	Tyr	Lys	Gly	Ile	Ala	
	850					855					860					
Ser	Asn	Tyr	Leu	Ala	Glu	Leu	Gln	Glu	Gly	Asp	Thr	Ile	Thr	Cys	Phe	
865					870					875					880	
Ile	Ser	Thr	Pro	Gln	Ser	Glu	Phe	Thr	Leu	Pro	Lys	Asp	Pro	Glu	Thr	
				885					890					895		
Pro	Leu	Ile	Met	Val	Gly	Pro	Gly	Thr	Gly	Val	Ala	Pro	Phe	Arg	Gly	

900					905					910					
Phe	Val	Gln	Ala	Arg	Lys	Gln	Leu	Lys	Glu	Gln	Gly	Gln	Ser	Leu	Gly
	915						920					925			
Glu	Ala	His	Leu	Tyr	Phe	Gly	Cys	Arg	Ser	Pro	His	Glu	Asp	Tyr	Leu
	930					935					940				
Tyr	Gln	Glu	Glu	Leu	Glu	Asn	Ala	Gln	Ser	Glu	Gly	Ile	Ile	Thr	Leu
945					950					955					960
His	Thr	Ala	Phe	Ser	Arg	Met	Pro	Asn	Gln	Pro	Lys	Thr	Tyr	Val	Gln
				965					970					975	
His	Val	Met	Glu	Gln	Asp	Gly	Lys	Lys	Leu	Ile	Glu	Leu	Leu	Asp	Gln
			980					985					990		
Gly	Ala	His	Phe	Tyr	Ile	Cys	Gly	Asp	Gly	Ser	Gln	Met	Ala	Pro	Ala
		995					1000					1005			
Val	Glu	Ala	Thr	Leu	Met	Lys	Ser	Tyr	Ala	Asp	Val	His	Gln	Val	Ser
	1010					1015					1020				
Glu	Ala	Asp	Ala	Arg	Leu	Trp	Leu	Gln	Gln	Leu	Glu	Glu	Lys	Gly	Arg
1025					1030				1035						1040
Tyr	Ala	Lys	Asp	Val	Trp	Ala	Gly								
				1045											

<210> 3
 <211> 30
 <212> DNA
 <213> Synthetic sequence

<220>
 <221> unsure
 <222> 1..30
 <223> n is g, a, t or c.

<220>
 <223> Description of the synthetic sequence: PCR primer

<400> 3

gcaggagacg gggtggnnac aagctggacg

30

<210> 4
 <211> 30
 <212> DNA
 <213> Synthetic sequence

<220>
 <221> unsure
 <222> 1..30
 <223> n is g, a, t or c.

<223> Description of the synthetic sequence: PCR primer

<400> 4

cgccagctt gtinncaacc cgtctctgc 30

<210> 5
<211> 34
<212> DNA
<213> Synthetic sequence

<220>
<221> unsure
<222> 1..34
<223> n is g, a, t or c.

<223> Description of the synthetic sequence: PCR primer

<400> 5
gaagcaatga acaagnnca gcgagcaaatt ccag 34

<210> 6
<211> 30
<212> DNA
<213> Synthetic sequence

<220>
<221> unsure
<222> 1..30
<223> n is g, a, t or c.

<223> Description of the synthetic sequence: PCR primer

<400> 6
ctggatttgc tcgctgnnnc ttgttcattg 30

<210> 7
<211> 41
<212> DNA
<213> Synthetic sequence

<220>
<221> unsure
<222> 1..41
<223> n is g, a, t or c.

<223> Description of the synthetic sequence: PCR primer

<400> 7
gctttgataa aaacttaaag tcaannnctt aaatttgtac g 41

<210> 8
<211> 40
<212> DNA
<213> Synthetic sequence

<220>
<221> unsure
<222> 1..40
<223> n is g, a, t or c.

<223> Description of the synthetic sequence: PCR primer

11

<400> 8

cgtacaaatt taagnnnttg acttaagttt ttatcaaagc

40

<210> 9

<211> 1049

<212> PRT

<213> Bacillus megaterium

<400> 9

Met	Thr	Ile	Lys	Glu	Met	Pro	Gln	Pro	Lys	Thr	Phe	Gly	Glu	Leu	Lys	
1				5					10					15		
Asn	Leu	Pro	Leu	Leu	Asn	Thr	Asp	Lys	Pro	Val	Gln	Ala	Leu	Met	Lys	
			20					25					30			
Ile	Ala	Asp	Glu	Leu	Gly	Glu	Ile	Phe	Lys	Phe	Glu	Ala	Pro	Gly	Arg	
		35					40					45				
Val	Thr	Arg	Tyr	Leu	Ser	Ser	Gln	Arg	Leu	Ile	Lys	Glu	Ala	Cys	Asp	
	50					55					60					
Glu	Ser	Arg	Phe	Asp	Lys	Asn	Leu	Ser	Gln	Ala	Leu	Lys	Phe	Val	Arg	
65					70					75					80	
Asp	Phe	Ala	Gly	Asp	Gly	Leu	Phe	Thr	Ser	Trp	Thr	His	Glu	Lys	Asn	
			85						90					95		
Trp	Lys	Lys	Ala	His	Asn	Ile	Leu	Leu	Pro	Ser	Phe	Ser	Gln	Gln	Ala	
			100					105						110		
Met	Lys	Gly	Tyr	His	Ala	Met	Met	Val	Asp	Ile	Ala	Val	Gln	Leu	Val	
		115					120					125				
Gln	Lys	Trp	Glu	Arg	Leu	Asn	Ala	Asp	Glu	His	Ile	Glu	Val	Pro	Glu	
		130				135					140					
Asp	Met	Thr	Arg	Leu	Thr	Leu	Asp	Thr	Ile	Gly	Leu	Cys	Gly	Phe	Asn	
145					150					155					160	
Tyr	Arg	Phe	Asn	Ser	Phe	Tyr	Arg	Asp	Gln	Pro	His	Pro	Phe	Ile	Thr	
			165						170					175		
Ser	Met	Val	Arg	Ala	Leu	Asp	Glu	Ala	Met	Asn	Lys	Leu	Gln	Arg	Ala	
			180					185					190			
Asn	Pro	Asp	Asp	Pro	Ala	Tyr	Asp	Glu	Asn	Lys	Arg	Gln	Phe	Gln	Glu	
		195					200					205				
Asp	Ile	Lys	Val	Met	Asn	Asp	Leu	Val	Asp	Lys	Ile	Ile	Ala	Asp	Arg	
	210					215					220					
Lys	Ala	Ser	Gly	Glu	Gln	Ser	Asp	Asp	Leu	Leu	Thr	His	Met	Leu	Asn	
225					230					235					240	
Gly	Lys	Asp	Pro	Glu	Thr	Gly	Glu	Pro	Leu	Asp	Asp	Glu	Asn	Ile	Arg	
				245					250					255		

12

Tyr Gln Ile Ile Thr Phe Leu Ile Ala Gly His Glu Thr Thr Ser Gly
 260 265 270
 Leu Leu Ser Phe Ala Leu Tyr Phe Leu Val Lys Asn Pro His Val Leu
 275 280 285
 Gln Lys Ala Ala Glu Glu Ala Ala Arg Val Leu Val Asp Pro Val Pro
 290 295 300
 Ser Tyr Lys Gln Val Lys Gln Leu Lys Tyr Val Gly Met Val Leu Asn
 305 310 315 320
 Glu Ala Leu Arg Leu Trp Pro Thr Ala Pro Ala Phe Ser Leu Tyr Ala
 325 330 335
 Lys Glu Asp Thr Val Leu Gly Gly Glu Tyr Pro Leu Glu Lys Gly Asp
 340 345 350
 Glu Leu Met Val Leu Ile Pro Gln Leu His Arg Asp Lys Thr Ile Trp
 355 360 365
 Gly Asp Asp Val Glu Glu Phe Arg Pro Glu Arg Phe Glu Asn Pro Ser
 370 375 380
 Ala Ile Pro Gln His Ala Phe Lys Pro Phe Gly Asn Gly Gln Arg Ala
 385 390 395 400
 Cys Ile Gly Gln Gln Phe Ala Leu His Glu Ala Thr Leu Val Leu Gly
 405 410 415
 Met Met Leu Lys His Phe Asp Phe Glu Asp His Thr Asn Tyr Glu Leu
 420 425 430
 Asp Ile Lys Glu Thr Leu Thr Leu Lys Pro Glu Gly Phe Val Val Lys
 435 440 445
 Ala Lys Ser Lys Lys Ile Pro Leu Gly Gly Ile Pro Ser Pro Ser Thr
 450 455 460
 Glu Gln Ser Ala Lys Lys Val Arg Lys Lys Ala Glu Asn Ala His Asn
 465 470 475 480
 Thr Pro Leu Leu Val Leu Tyr Gly Ser Asn Met Gly Thr Ala Glu Gly
 485 490 495
 Thr Ala Arg Asp Leu Ala Asp Ile Ala Met Ser Lys Gly Phe Ala Pro
 500 505 510
 Gln Val Ala Thr Leu Asp Ser His Ala Gly Asn Leu Pro Arg Glu Gly
 515 520 525
 Ala Val Leu Ile Val Thr Ala Ser Tyr Asn Gly His Pro Pro Asp Asn
 530 535 540
 Ala Lys Gln Phe Val Asp Trp Leu Asp Gln Ala Ser Ala Asp Glu Val
 545 550 555 560
 Lys Gly Val Arg Tyr Ser Val Phe Gly Cys Gly Asp Lys Asn Trp Ala
 565 570 575

Thr Thr Tyr Gln Lys Val Pro Ala Phe Ile Asp Glu Thr Leu Ala Ala
 580 585 590
 Lys Gly Ala Glu Asn Ile Ala Asp Arg Gly Glu Ala Asp Ala Ser Asp
 595 600 605
 Asp Phe Glu Gly Thr Tyr Glu Glu Trp Arg Glu His Met Trp Ser Asp
 610 615 620
 Val Ala Ala Tyr Phe Asn Leu Asp Ile Glu Asn Ser Glu Asp Asn Lys
 625 630 635 640
 Ser Thr Leu Ser Leu Gln Phe Val Asp Ser Ala Ala Asp Met Pro Leu
 645 650 655
 Ala Lys Met His Gly Ala Phe Ser Thr Asn Val Val Ala Ser Lys Glu
 660 665 670
 Leu Gln Gln Pro Gly Ser Ala Arg Ser Thr Arg His Leu Glu Ile Glu
 675 680 685
 Leu Pro Lys Glu Ala Ser Tyr Gln Glu Gly Asp His Leu Gly Val Ile
 690 695 700
 Pro Arg Asn Tyr Glu Gly Ile Val Asn Arg Val Thr Ala Arg Phe Gly
 705 710 715 720
 Leu Asp Ala Ser Gln Gln Ile Arg Leu Glu Ala Glu Glu Glu Lys Leu
 725 730 735
 Ala His Leu Pro Leu Ala Lys Thr Val Ser Val Glu Glu Leu Leu Gln
 740 745 750
 Tyr Val Glu Leu Gln Asp Pro Val Thr Arg Thr Gln Leu Arg Ala Met
 755 760 765
 Ala Ala Lys Thr Val Cys Pro Pro His Lys Val Glu Leu Glu Ala Leu
 770 775 780
 Leu Glu Lys Gln Ala Tyr Lys Glu Gln Val Leu Ala Lys Arg Leu Thr
 785 790 795 800
 Met Leu Glu Leu Leu Glu Lys Tyr Pro Ala Cys Glu Met Lys Phe Ser
 805 810 815
 Glu Phe Ile Ala Leu Leu Pro Ser Ile Arg Pro Arg Tyr Tyr Ser Ile
 820 825 830
 Ser Ser Ser Pro Arg Val Asp Glu Lys Gln Ala Ser Ile Thr Val Ser
 835 840 845
 Val Val Ser Gly Glu Ala Trp Ser Gly Tyr Gly Glu Tyr Lys Gly Ile
 850 855 860
 Ala Ser Asn Tyr Leu Ala Glu Leu Gln Glu Gly Asp Thr Ile Thr Cys
 865 870 875 880
 Phe Ile Ser Thr Pro Gln Ser Glu Phe Thr Leu Pro Lys Asp Pro Glu

14

885	890	895
Thr Pro Leu Ile Met Val Gly Pro Gly Thr Gly Val Ala Pro Phe Arg		
900	905	910
Gly Phe Val Gln Ala Arg Lys Gln Leu Lys Glu Gln Gly Gln Ser Leu		
915	920	925
Gly Glu Ala His Leu Tyr Phe Gly Cys Arg Ser Pro His Glu Asp Tyr		
930	935	940
Leu Tyr Gln Glu Glu Leu Glu Asn Ala Gln Ser Glu Gly Ile Ile Thr		
945	950	955
Leu His Thr Ala Phe Ser Arg Met Pro Asn Gln Pro Lys Thr Tyr Val		
965	970	975
Gln His Val Met Glu Gln Asp Gly Lys Lys Leu Ile Glu Leu Leu Asp		
980	985	990
Gln Gly Ala His Phe Tyr Ile Cys Gly Asp Gly Ser Gln Met Ala Pro		
995	1000	1005
Ala Val Glu Ala Thr Leu Met Lys Ser Tyr Ala Asp Val His Gln Val		
1010	1015	1020
Ser Glu Ala Asp Ala Arg Leu Trp Leu Gln Gln Leu Glu Glu Lys Gly		
1025	1030	1035
Arg Tyr Ala Lys Asp Val Trp Ala Gly		
1045		1040

0050/50915

26

gat ttt gca gga gac ggg tta ttt aca agc tgg acg cat gaa aaa aat	288
Asp Phe Ala Gly Asp Gly Leu Phe Thr Ser Trp Thr His Glu Lys Asn	
80 85 90 95	
tgg aaa aaa gcg cat aat atc tta ctt cca agc ttc agt cag cag gca	336
Trp Lys Lys Ala His Asn Ile Leu Leu Pro Ser Phe Ser Gln Gln Ala	
100 105 110	
atg aaa ggc tat cat gcg atg atg gtc gat atc gcc gtg cag ctt gtt	384
Met Lys Gly Tyr His Ala Met Met Val Asp Ile Ala Val Gln Leu Val	
115 120 125	
caa aag tgg gag cgt cta aat gca gat gag cat att gaa gta ccg gaa	432
Gln Lys Trp Glu Arg Leu Asn Ala Asp Glu His Ile Glu Val Pro Glu	
130 135 140	
gac atg aca cgt tta acg ctt gat aca att ggt ctt tgc ggc ttt aac	480
Asp Met Thr Arg Leu Thr Leu Asp Thr Ile Gly Leu Cys Gly Phe Asn	
145 150 155	
tat cgc ttt aac agc ttt tac cga gat cag cct cat cca ttt att aca	528
Tyr Arg Phe Asn Ser Phe Tyr Arg Asp Gln Pro His Pro Phe Ile Thr	
160 165 170 175	
agt atg gtc cgt gca ctg gat gaa gca atg aac aag ctg cag cga gca	576
Ser Met Val Arg Ala Leu Asp Glu Ala Met Asn Lys Leu Gln Arg Ala	
180 185 190	
aat cca gac gac cca gct tat gat gaa aac aag cgc cag ttt caa gaa	624
Asn Pro Asp Asp Pro Ala Tyr Asp Glu Asn Lys Arg Gln Phe Gln Glu	
195 200 205	
gat atc aag gtg atg aac gac cta gta gat aaa att att gca gat cgc	672
Asp Ile Lys Val Met Asn Asp Leu Val Asp Lys Ile Ile Ala Asp Arg	
210 215 220	
aaa gca agc ggt gaa caa agc gat gat tta tta acg cat atg cta aac	720
Lys Ala Ser Gly Glu Gln Ser Asp Asp Leu Leu Thr His Met Leu Asn	
225 230 235	
gga aaa gat cca gaa acg ggt gag ccg ctt gat gac gag aac att cgc	768
Gly Lys Asp Pro Glu Thr Gly Glu Pro Leu Asp Asp Glu Asn Ile Arg	
240 245 250 255	
tat caa att att aca ttc tta att gcg gga cac gaa aca aca agt ggt	816
Tyr Gln Ile Ile Thr Phe Leu Ile Ala Gly His Glu Thr Thr Ser Gly	
260 265 270	
ctt tta tca ttt gcg ctg tat ttc tta gtg aaa aat cca cat gta tta	864
Leu Leu Ser Phe Ala Leu Tyr Phe Leu Val Lys Asn Pro His Val Leu	
275 280 285	

27

caa aaa gca gca gaa gaa gca gca cga gtt cta gta gat cct gtt cca	912
Gln Lys Ala Ala Glu Glu Ala Ala Arg Val Leu Val Asp Pro Val Pro	
290 295 300	
agc tac aaa caa gtc aaa cag ctt aaa tat gtc ggc atg gtc tta aac	960
Ser Tyr Lys Lys Val Lys Lys Leu Lys Tyr Val Gly Met Val Leu Asn	
305 310 315	
gaa gcg ctg cgc tta tgg cca act gct cct gcg ttt tcc cta tat gca	1008
Glu Ala Leu Arg Leu Trp Pro Thr Ala Pro Ala Phe Ser Leu Tyr Ala	
320 325 330 335	
aaa gaa gat acg gtg ctt gga gga gaa tat cct tta gaa aaa ggc gac	1056
Lys Glu Asp Thr Val Leu Gly Gly Glu Tyr Pro Leu Glu Lys Gly Asp	
340 345 350	
gaa cta atg gtt ctg att cct cag ctt cac cgt gat aaa aca att tgg	1104
Glu Leu Met Val Leu Ile Pro Gln Leu His Arg Asp Lys Thr Ile Trp	
355 360 365	
gga gac gat gtg gaa gag ttc cgt cca gag cgt ttt gaa aat cca agt	1152
Gly Asp Asp Val Glu Glu Phe Arg Pro Glu Arg Phe Glu Asn Pro Ser	
370 375 380	
gcg att ccg cag cat gcg ttt aaa ccg ttt gga aac ggt cag cgt gcg	1200
Ala Ile Pro Gln His Ala Phe Lys Pro Phe Gly Asn Gly Gln Arg Ala	
385 390 395	
tgt atc ggt cag cag ttc gct ctt cat gaa gca acg ctg gta ctt ggt	1248
Cys Ile Gly Gln Gln Phe Ala Leu His Glu Ala Thr Leu Val Leu Gly	
400 405 410 415	
atg atg cta aaa cac ttt gac ttt gaa gat cat aca aac tac gag ctg	1296
Met Met Leu Lys His Phe Asp Phe Glu Asp His Thr Asn Tyr Glu Leu	
420 425 430	
gat att aaa gaa act tta acg tta aaa cct gaa ggc ttt gtg gta aaa	1344
Asp Ile Lys Glu Thr Leu Thr Leu Lys Pro Glu Gly Phe Val Val Lys	
435 440 445	
gca aaa tcg aaa aaa att ccg ctt ggc ggt att cct tca cct agc act	1392
Ala Lys Ser Lys Lys Ile Pro Leu Gly Gly Ile Pro Ser Pro Ser Thr	
450 455 460	
gaa cag tct gct aaa aaa gta cgc aaa aag gca gaa aac gct cat aat	1440
Glu Gln Ser Ala Lys Lys Val Arg Lys Lys Ala Glu Asn Ala His Asn	
465 470 475	
acg ccg ctg ctt gtg cta tac ggt tca aat atg gga aca gct gaa gga	1488
Thr Pro Leu Leu Val Leu Tyr Gly Ser Asn Met Gly Thr Ala Glu Gly	
480 485 490 495	

0050/50915

28

acg gcg cgt gat tta gca gat att gca atg agc aaa gga ttt gca ccg	1536
Thr Ala Arg Asp Leu Ala Asp Ile Ala Met Ser Lys Gly Phe Ala Pro	
500 505 510	
cag gtc gca acg ctt gat tca cac gcc gga aat ctt ccg cgc gaa gga	1584
Gln Val Ala Thr Leu Asp Ser His Ala Gly Asn Leu Pro Arg Glu Gly	
515 520 525	
gct gta tta att gta acg gcg tct tat aac ggt cat ccg cct gat aac	1632
Ala Val Leu Ile Val Thr Ala Ser Tyr Asn Gly His Pro Pro Asp Asn	
530 535 540	
gca aag caa ttt gtc gac tgg tta gac caa gcg tct gct gat gaa gta	1680
Ala Lys Gln Phe Val Asp Trp Leu Asp Gln Ala Ser Ala Asp Glu Val	
545 550 555	
aaa ggc gtt cgc tac tcc gta ttt gga tgc ggc gat aaa aac tgg gct	1728
Lys Gly Val Arg Tyr Ser Val Phe Gly Cys Gly Asp Lys Asn Trp Ala	
560 565 570 575	
act acg tat caa aaa gtg cct gct ttt atc gat gaa acg ctt gcc gct	1776
Thr Thr Tyr Gln Lys Val Pro Ala Phe Ile Asp Glu Thr Leu Ala Ala	
580 585 590	
aaa ggg gca gaa aac atc gct gac cgc ggt gaa gca gat gca agc gac	1824
Lys Gly Ala Glu Asn Ile Ala Asp Arg Gly Glu Ala Asp Ala Ser Asp	
595 600 605	
gac ttt gaa ggc aca tat gaa gaa tgg cgt gaa cat atg tgg agt gac	1872
Asp Phe Glu Gly Thr Tyr Glu Glu Trp Arg Glu His Met Trp Ser Asp	
610 615 620	
gta gca gcc tac ttt aac ctc gac att gaa aac agt gaa gat aat aaa	1920
Val Ala Ala Tyr Phe Asn Leu Asp Ile Glu Asn Ser Glu Asp Asn Lys	
625 630 635	
tct act ctt tca ctt caa ttt gtc gac agc gcc gcg gat atg ccg ctt	1968
Ser Thr Leu Ser Leu Gln Phe Val Asp Ser Ala Ala Asp Met Pro Leu	
640 645 650 655	
gcg aaa atg cac ggt gcg ttt tca acg aac gtc gta gca agc aaa gaa	2016
Ala Lys Met His Gly Ala Phe Ser Thr Asn Val Val Ala Ser Lys Glu	
660 665 670	
ctt caa cag cca ggc agt gca cga agc acg cga cat ctt gaa att gaa	2064
Leu Gln Gln Pro Gly Ser Ala Arg Ser Thr Arg His Leu Glu Ile Glu	
675 680 685	
ctt cca aaa gaa gct tct tat caa gaa gga gat cat tta ggt gtt att	2112
Leu Pro Lys Glu Ala Ser Tyr Gln Glu Gly Asp His Leu Gly Val Ile	
690 695 700	

29

cct cgc aac tat gaa gga ata gta aac cgt gta aca gca agg ttc ggc	2160
Pro Arg Asn Tyr Glu Gly Ile Val Asn Arg Val Thr Ala Arg Phe Gly	
705 710 715	
cta gat gca tca cag caa atc cgt ctg gaa gca gaa gaa gaa aaa tta	2208
Leu Asp Ala Ser Gln Gln Ile Arg Leu Glu Ala Glu Glu Glu Lys Leu	
720 725 730 735	
gct cat ttg cca ctc gct aaa aca gta tcc gta gaa gag ctt ctg caa	2256
Ala His Leu Pro Leu Ala Lys Thr Val Ser Val Glu Glu Leu Leu Gln	
740 745 750	
tac gtg gag ctt caa gat cct gtt acg cgc acg cag ctt cgc gca atg	2304
Tyr Val Glu Leu Gln Asp Pro Val Thr Arg Thr Gln Leu Arg Ala Met	
755 760 765	
gct gct aaa acg gtc tgc ccg ccg cat aaa gta gag ctt gaa gcc ttg	2352
Ala Ala Lys Thr Val Cys Pro Pro His Lys Val Glu Leu Glu Ala Leu	
770 775 780	
ctt gaa aag caa gcc tac aaa gaa caa gtg ctg gca aaa cgt tta aca	2400
Leu Glu Lys Gln Ala Tyr Lys Glu Gln Val Leu Ala Lys Arg Leu Thr	
785 790 795	
atg ctt gaa ctg ctt gaa aaa tac ccg gcg tgt gaa atg aaa ttc agc	2448
Met Leu Glu Leu Leu Glu Lys Tyr Pro Ala Cys Glu Met Lys Phe Ser	
800 805 810 815	
gaa ttt atc gcc ctt ctg cca agc ata cgc ccg cgc tat tac tcg att	2496
Glu Phe Ile Ala Leu Leu Pro Ser Ile Arg Pro Arg Tyr Tyr Ser Ile	
820 825 830	
tct tca tca cct cgt gtc gat gaa aaa caa gca agc atc acg gtc agc	2544
Ser Ser Ser Pro Arg Val Asp Glu Lys Gln Ala Ser Ile Thr Val Ser	
835 840 845	
gtt gtc tca gga gaa gcg tgg agc gga tat gga gaa tat aaa gga att	2592
Val Val Ser Gly Glu Ala Trp Ser Gly Tyr Gly Glu Tyr Lys Gly Ile	
850 855 860	
gcg tcg aac tat ctt gcc gag ctg caa gaa gga gat acg att acg tgc	2640
Ala Ser Asn Tyr Leu Ala Glu Leu Gln Glu Gly Asp Thr Ile Thr Cys	
865 870 875	
ttt att tcc aca ccg cag tca gaa ttt acg ctg cca aaa gac cct gaa	2688
Phe Ile Ser Thr Pro Gln Ser Glu Phe Thr Leu Pro Lys Asp Pro Glu	
880 885 890 895	
acg ccg ctt atc atg gtc gga ccg gga aca ggc gtc gcg ccg ttt aga	2736
Thr Pro Leu Ile Met Val Gly Pro Gly Thr Gly Val Ala Pro Phe Arg	
900 905 910	

0050/50915

30

ggc ttt gtg cag gcg cgc aaa cag cta aaa gaa caa gga cag tca ctt 2784
 Gly Phe Val Gln Ala Arg Lys Gln Leu Lys Glu Gln Gly Gln Ser Leu
 915 920 925

gga gaa gca cat tta tac ttc ggc tgc cgt tca cct cat gaa gac tat 2832
 Gly Glu Ala His Leu Tyr Phe Gly Cys Arg Ser Pro His Glu Asp Tyr
 930 935 940

ctg tat caa gaa gag ctt gaa aac gcc caa agc gaa ggc atc att acg 2880
 Leu Tyr Gln Glu Glu Leu Glu Asn Ala Gln Ser Glu Gly Ile Ile Thr
 945 950 955

ctt cat acc gct ttt tct cgc atg cca aat cag ccg aaa aca tac gtt 2928
 Leu His Thr Ala Phe Ser Arg Met Pro Asn Gln Pro Lys Thr Tyr Val
 960 965 970 975

cag cac gta atg gaa caa gac ggc aag aaa ttg att gaa ctt ctt gat 2976
 Gln His Val Met Glu Gln Asp Gly Lys Lys Leu Ile Glu Leu Leu Asp
 980 985 990

caa gga gcg cac ttc tat att tgc gga gac gga agc caa atg gca cct 3024
 Gln Gly Ala His Phe Tyr Ile Cys Gly Asp Gly Ser Gln Met Ala Pro
 995 1000 1005

gcc gtt gaa gca acg ctt atg aaa agc tat gct gac gtt cac caa gtg 3072
 Ala Val Glu Ala Thr Leu Met Lys Ser Tyr Ala Asp Val His Gln Val
 1010 1015 1020

agt gaa gca gac gct cgc tta tgg ctg cag cag cta gaa gaa aaa ggc 3120
 Ser Glu Ala Asp Ala Arg Leu Trp Leu Gln Gln Leu Glu Glu Lys Gly
 1025 1030 1035

cga tac gca aaa gac gtg tgg gct ggg taa 3150
 Arg Tyr Ala Lys Asp Val Trp Ala Gly
 1040 1045

<210> 2

<211> 1048

<212> PRT

<213> Bacillus megaterium

<400> 2

Thr Ile Lys Glu Met Pro Gln Pro Lys Thr Phe Gly Glu Leu Lys Asn
 1 5 10 15

Leu Pro Leu Leu Asn Thr Asp Lys Pro Val Gln Ala Leu Met Lys Ile
 20 25 30

Ala Asp Glu Leu Gly Glu Ile Phe Lys Phe Glu Ala Pro Gly Arg Val
 35 40 45

0050/50915

31

Thr	Arg	Tyr	Leu	Ser	Ser	Gln	Arg	Leu	Ile	Lys	Glu	Ala	Cys	Asp	Glu
50						55					60				
Ser	Arg	Phe	Asp	Lys	Asn	Leu	Ser	Gln	Ala	Leu	Lys	Phe	Val	Arg	Asp
65					70					75					80
Phe	Ala	Gly	Asp	Gly	Leu	Phe	Thr	Ser	Trp	Thr	His	Glu	Lys	Asn	Trp
				85					90					95	
Lys	Lys	Ala	His	Asn	Ile	Leu	Leu	Pro	Ser	Phe	Ser	Gln	Gln	Ala	Met
			100					105					110		
Lys	Gly	Tyr	His	Ala	Met	Met	Val	Asp	Ile	Ala	Val	Gln	Leu	Val	Gln
	115						120					125			
Lys	Trp	Glu	Arg	Leu	Asn	Ala	Asp	Glu	His	Ile	Glu	Val	Pro	Glu	Asp
130						135					140				
Met	Thr	Arg	Leu	Thr	Leu	Asp	Thr	Ile	Gly	Leu	Cys	Gly	Phe	Asn	Tyr
145					150					155					160
Arg	Phe	Asn	Ser	Phe	Tyr	Arg	Asp	Gln	Pro	His	Pro	Phe	Ile	Thr	Ser
				165					170					175	
Met	Val	Arg	Ala	Leu	Asp	Glu	Ala	Met	Asn	Lys	Leu	Gln	Arg	Ala	Asn
			180					185					190		
Pro	Asp	Asp	Pro	Ala	Tyr	Asp	Glu	Asn	Lys	Arg	Gln	Phe	Gln	Glu	Asp
	195						200					205			
Ile	Lys	Val	Met	Asn	Asp	Leu	Val	Asp	Lys	Ile	Ile	Ala	Asp	Arg	Lys
210						215					220				
Ala	Ser	Gly	Glu	Gln	Ser	Asp	Asp	Leu	Leu	Thr	His	Met	Leu	Asn	Gly
225					230					235					240
Lys	Asp	Pro	Glu	Thr	Gly	Glu	Pro	Leu	Asp	Asp	Glu	Asn	Ile	Arg	Tyr
				245					250					255	
Gln	Ile	Ile	Thr	Phe	Leu	Ile	Ala	Gly	His	Glu	Thr	Thr	Ser	Gly	Leu
			260					265					270		
Leu	Ser	Phe	Ala	Leu	Tyr	Phe	Leu	Val	Lys	Asn	Pro	His	Val	Leu	Gln
		275					280					285			
Lys	Ala	Ala	Glu	Glu	Ala	Ala	Arg	Val	Leu	Val	Asp	Pro	Val	Pro	Ser
	290					295					300				
Tyr	Lys	Gln	Val	Lys	Gln	Leu	Lys	Tyr	Val	Gly	Met	Val	Leu	Asn	Glu
305					310					315				320	

0050/50915

32

Ala Leu Arg Leu Trp Pro Thr Ala Pro Ala Phe Ser Leu Tyr Ala Lys
325 330 335

Glu Asp Thr Val Leu Gly Gly Glu Tyr Pro Leu Glu Lys Gly Asp Glu
340 345 350

Leu Met Val Leu Ile Pro Gln Leu His Arg Asp Lys Thr Ile Trp Gly
355 360 365

Asp Asp Val Glu Glu Phe Arg Pro Glu Arg Phe Glu Asn Pro Ser Ala
370 375 380

Ile Pro Gln His Ala Phe Lys Pro Phe Gly Asn Gly Gln Arg Ala Cys
385 390 395 400

Ile Gly Gln Gln Phe Ala Leu His Glu Ala Thr Leu Val Leu Gly Met
405 410 415

Met Leu Lys His Phe Asp Phe Glu Asp His Thr Asn Tyr Glu Leu Asp
420 425 430

Ile Lys Glu Thr Leu Thr Leu Lys Pro Glu Gly Phe Val Val Lys Ala
435 440 445

Lys Ser Lys Lys Ile Pro Leu Gly Gly Ile Pro Ser Pro Ser Thr Glu
450 455 460

Gln Ser Ala Lys Lys Val Arg Lys Lys Ala Glu Asn Ala His Asn Thr
465 470 475 480

Pro Leu Leu Val Leu Tyr Gly Ser Asn Met Gly Thr Ala Glu Gly Thr
485 490 495

Ala Arg Asp Leu Ala Asp Ile Ala Met Ser Lys Gly Phe Ala Pro Gln
500 505 510

Val Ala Thr Leu Asp Ser His Ala Gly Asn Leu Pro Arg Glu Gly Ala
515 520 525

Val Leu Ile Val Thr Ala Ser Tyr Asn Gly His Pro Pro Asp Asn Ala
530 535 540

Lys Gln Phe Val Asp Trp Leu Asp Gln Ala Ser Ala Asp Glu Val Lys
545 550 555 560

Gly Val Arg Tyr Ser Val Phe Gly Cys Gly Asp Lys Asn Trp Ala Thr
565 570 575

Thr Tyr Gln Lys Val Pro Ala Phe Ile Asp Glu Thr Leu Ala Ala Lys
580 585 590

0050/50915

33

Gly	Ala	Glu	Asn	Ile	Ala	Asp	Arg	Gly	Glu	Ala	Asp	Ala	Ser	Asp	Asp	595	600	605	
Phe	Glu	Gly	Thr	Tyr	Glu	Glu	Trp	Arg	Glu	His	Met	Trp	Ser	Asp	Val	610	615	620	
Ala	Ala	Tyr	Phe	Asn	Leu	Asp	Ile	Glu	Asn	Ser	Glu	Asp	Asn	Lys	Ser	625	630	635	640
Thr	Leu	Ser	Leu	Gln	Phe	Val	Asp	Ser	Ala	Ala	Asp	Met	Pro	Leu	Ala	645	650	655	
Lys	Met	His	Gly	Ala	Phe	Ser	Thr	Asn	Val	Val	Ala	Ser	Lys	Glu	Leu	660	665	670	
Gln	Gln	Pro	Gly	Ser	Ala	Arg	Ser	Thr	Arg	His	Leu	Glu	Ile	Glu	Leu	675	680	685	
Pro	Lys	Glu	Ala	Ser	Tyr	Gln	Glu	Gly	Asp	His	Leu	Gly	Val	Ile	Pro	690	695	700	
Arg	Asn	Tyr	Glu	Gly	Ile	Val	Asn	Arg	Val	Thr	Ala	Arg	Phe	Gly	Leu	705	710	715	720
Asp	Ala	Ser	Gln	Gln	Ile	Arg	Leu	Glu	Ala	Glu	Glu	Glu	Lys	Leu	Ala	725	730	735	
His	Leu	Pro	Leu	Ala	Lys	Thr	Val	Ser	Val	Glu	Glu	Leu	Leu	Gln	Tyr	740	745	750	
Val	Glu	Leu	Gln	Asp	Pro	Val	Thr	Arg	Thr	Gln	Leu	Arg	Ala	Met	Ala	755	760	765	
Ala	Lys	Thr	Val	Cys	Pro	Pro	His	Lys	Val	Glu	Leu	Glu	Ala	Leu	Leu	770	775	780	
Glu	Lys	Gln	Ala	Tyr	Lys	Glu	Gln	Val	Leu	Ala	Lys	Arg	Leu	Thr	Met	785	790	795	800
Leu	Glu	Leu	Leu	Glu	Lys	Tyr	Pro	Ala	Cys	Glu	Met	Lys	Phe	Ser	Glu	805	810	815	
Phe	Ile	Ala	Leu	Leu	Pro	Ser	Ile	Arg	Pro	Arg	Tyr	Tyr	Ser	Ile	Ser	820	825	830	
Ser	Ser	Pro	Arg	Val	Asp	Glu	Lys	Gln	Ala	Ser	Ile	Thr	Val	Ser	Val	835	840	845	
Val	Ser	Gly	Glu	Ala	Trp	Ser	Gly	Tyr	Gly	Glu	Tyr	Lys	Gly	Ile	Ala	850	855	860	

0050/50915

34

Ser Asn Tyr Leu Ala Glu Leu Gln Glu Gly Asp Thr Ile Thr Cys Phe
 865 870 875 880
 Ile Ser Thr Pro Gln Ser Glu Phe Thr Leu Pro Lys Asp Pro Glu Thr
 885 890 895
 Pro Leu Ile Met Val Gly Pro Gly Thr Gly Val Ala Pro Phe Arg Gly
 900 905 910
 Phe Val Gln Ala Arg Lys Gln Leu Lys Glu Gln Gly Gln Ser Leu Gly
 915 920 925
 Glu Ala His Leu Tyr Phe Gly Cys Arg Ser Pro His Glu Asp Tyr Leu
 930 935 940
 Tyr Gln Glu Glu Leu Glu Asn Ala Gln Ser Glu Gly Ile Ile Thr Leu
 945 950 955 960
 His Thr Ala Phe Ser Arg Met Pro Asn Gln Pro Lys Thr Tyr Val Gln
 965 970 975
 His Val Met Glu Gln Asp Gly Lys Lys Leu Ile Glu Leu Leu Asp Gln
 980 985 990
 Gly Ala His Phe Tyr Ile Cys Gly Asp Gly Ser Gln Met Ala Pro Ala
 995 1000 1005
 Val Glu Ala Thr Leu Met Lys Ser Tyr Ala Asp Val His Gln Val Ser
 1010 1015 1020
 Glu Ala Asp Ala Arg Leu Trp Leu Gln Gln Leu Glu Glu Lys Gly Arg
 1025 1030 1035 1040
 Tyr Ala Lys Asp Val Trp Ala Gly
 1045

<210> 3

<211> 30

<212> DNA

<213> Synthetic sequence

<220>

<223> Description of the synthetic sequence: PCR primer

<400> 3

gcaggagacg ggttgnnnac aagctggacg

30

<210> 4

<211> 30

<212> DNA

<213> Synthetic sequence

0050/50915

35

<220>

<223> Description of the synthetic sequence: PCR primer

<400> 4

cgtccagctt gttnncaacc cgtctcctgc

30

<210> 5

<211> 34

<212> DNA

<213> Synthetic sequence

<220>

<223> Description of the synthetic sequence: PCR primer

<400> 5

gaagcaatga acaagnnnca gcgagcaaatt ccag

34

<210> 6

<211> 30

<212> DNA

<213> Synthetic sequence

<220>

<223> Description of the synthetic sequence: PCR primer

<400> 6

ctggatttgc tcgctgnnnc ttgttcattg

30

<210> 7

<211> 41

<212> DNA

<213> Synthetic sequence

<220>

<223> Description of the synthetic sequence: PCR primer

<400> 7

gctttgataa aaacttaaag tcaannnctt aaatttgtac g

41

<210> 8

<211> 40

<212> DNA

<213> Synthetic sequence

<220>

<223> Description of the synthetic sequence: PCR primer

<400> 8

cgtacaaatt taagnnnttg acttaagttt ttatcaaagc

40

0050/50915

36

<210> 9

<211> 1049

<212> PRT

<213> Bacillus megaterium

<400> 9

Met	Thr	Ile	Lys	Glu	Met	Pro	Gln	Pro	Lys	Thr	Phe	Gly	Glu	Leu	Lys
1				5					10					15	

Asn	Leu	Pro	Leu	Leu	Asn	Thr	Asp	Lys	Pro	Val	Gln	Ala	Leu	Met	Lys
			20					25					30		

Ile	Ala	Asp	Glu	Leu	Gly	Glu	Ile	Phe	Lys	Phe	Glu	Ala	Pro	Gly	Arg
		35					40					45			

Val	Thr	Arg	Tyr	Leu	Ser	Ser	Gln	Arg	Leu	Ile	Lys	Glu	Ala	Cys	Asp
	50					55					60				

Glu	Ser	Arg	Phe	Asp	Lys	Asn	Leu	Ser	Gln	Ala	Leu	Lys	Phe	Val	Arg
65					70					75					80

Asp	Phe	Ala	Gly	Asp	Gly	Leu	Phe	Thr	Ser	Trp	Thr	His	Glu	Lys	Asn
				85					90					95	

Trp	Lys	Lys	Ala	His	Asn	Ile	Leu	Leu	Pro	Ser	Phe	Ser	Gln	Gln	Ala
			100					105					110		

Met	Lys	Gly	Tyr	His	Ala	Met	Met	Val	Asp	Ile	Ala	Val	Gln	Leu	Val
		115					120					125			

Gln	Lys	Trp	Glu	Arg	Leu	Asn	Ala	Asp	Glu	His	Ile	Glu	Val	Pro	Glu
	130					135					140				

Asp	Met	Thr	Arg	Leu	Thr	Leu	Asp	Thr	Ile	Gly	Leu	Cys	Gly	Phe	Asn
145					150					155					160

Tyr	Arg	Phe	Asn	Ser	Phe	Tyr	Arg	Asp	Gln	Pro	His	Pro	Phe	Ile	Thr
			165						170					175	

Ser	Met	Val	Arg	Ala	Leu	Asp	Glu	Ala	Met	Asn	Lys	Leu	Gln	Arg	Ala
			180					185					190		

Asn	Pro	Asp	Asp	Pro	Ala	Tyr	Asp	Glu	Asn	Lys	Arg	Gln	Phe	Gln	Glu
		195					200					205			

Asp	Ile	Lys	Val	Met	Asn	Asp	Leu	Val	Asp	Lys	Ile	Ile	Ala	Asp	Arg
	210					215					220				

Lys	Ala	Ser	Gly	Glu	Gln	Ser	Asp	Asp	Leu	Leu	Thr	His	Met	Leu	Asn
225					230					235					240

Gly	Lys	Asp	Pro	Glu	Thr	Gly	Glu	Pro	Leu	Asp	Asp	Glu	Asn	Ile	Arg
				245						250		255			
Tyr	Gln	Ile	Ile	Thr	Phe	Leu	Ile	Ala	Gly	His	Glu	Thr	Thr	Ser	Gly
				260						265		270			
Leu	Leu	Ser	Phe	Ala	Leu	Tyr	Phe	Leu	Val	Lys	Asn	Pro	His	Val	Leu
				275						280		285			
Gln	Lys	Ala	Ala	Glu	Glu	Ala	Ala	Arg	Val	Leu	Val	Asp	Pro	Val	Pro
				290						295		300			
Ser	Tyr	Lys	Gln	Val	Lys	Gln	Leu	Lys	Tyr	Val	Gly	Met	Val	Leu	Asn
305				310				315				320			
Glu	Ala	Leu	Arg	Leu	Trp	Pro	Thr	Ala	Pro	Ala	Phe	Ser	Leu	Tyr	Ala
				325						330		335			
Lys	Glu	Asp	Thr	Val	Leu	Gly	Gly	Glu	Tyr	Pro	Leu	Glu	Lys	Gly	Asp
				340						345		350			
Glu	Leu	Met	Val	Leu	Ile	Pro	Gln	Leu	His	Arg	Asp	Lys	Thr	Ile	Trp
355				360				365							
Gly	Asp	Asp	Val	Glu	Glu	Phe	Arg	Pro	Glu	Arg	Phe	Glu	Asn	Pro	Ser
370				375				380							
Ala	Ile	Pro	Gln	His	Ala	Phe	Lys	Pro	Phe	Gly	Asn	Gly	Gln	Arg	Ala
385				390				395				400			
Cys	Ile	Gly	Gln	Gln	Phe	Ala	Leu	His	Glu	Ala	Thr	Leu	Val	Leu	Gly
				405						410		415			
Met	Met	Leu	Lys	His	Phe	Asp	Phe	Glu	Asp	His	Thr	Asn	Tyr	Glu	Leu
420				425				430							
Asp	Ile	Lys	Glu	Thr	Leu	Thr	Leu	Lys	Pro	Glu	Gly	Phe	Val	Val	Lys
435				440				445							
Ala	Lys	Ser	Lys	Lys	Ile	Pro	Leu	Gly	Gly	Ile	Pro	Ser	Pro	Ser	Thr
450				455				460							
Glu	Gln	Ser	Ala	Lys	Lys	Val	Arg	Lys	Lys	Ala	Glu	Asn	Ala	His	Asn
465				470				475				480			
Thr	Pro	Leu	Leu	Val	Leu	Tyr	Gly	Ser	Asn	Met	Gly	Thr	Ala	Glu	Gly
				485						490		495			
Thr	Ala	Arg	Asp	Leu	Ala	Asp	Ile	Ala	Met	Ser	Lys	Gly	Phe	Ala	Pro
500				505				510							

Gln	Val	Ala	Thr	Leu	Asp	Ser	His	Ala	Gly	Asn	Leu	Pro	Arg	Glu	Gly
515						520						525			
Ala	Val	Leu	Ile	Val	Thr	Ala	Ser	Tyr	Asn	Gly	His	Pro	Pro	Asp	Asn
530						535						540			
Ala	Lys	Gln	Phe	Val	Asp	Trp	Leu	Asp	Gln	Ala	Ser	Ala	Asp	Glu	Val
545						550						555			
Lys	Gly	Val	Arg	Tyr	Ser	Val	Phe	Gly	Cys	Gly	Asp	Lys	Asn	Trp	Ala
			565						570			575			
Thr	Thr	Tyr	Gln	Lys	Val	Pro	Ala	Phe	Ile	Asp	Glu	Thr	Leu	Ala	Ala
			580						585			590			
Lys	Gly	Ala	Glu	Asn	Ile	Ala	Asp	Arg	Gly	Glu	Ala	Asp	Ala	Ser	Asp
595						600						605			
Asp	Phe	Glu	Gly	Thr	Tyr	Glu	Glu	Trp	Arg	Glu	His	Met	Trp	Ser	Asp
610						615						620			
Val	Ala	Ala	Tyr	Phe	Asn	Leu	Asp	Ile	Glu	Asn	Ser	Glu	Asp	Asn	Lys
625						630						635			
Ser	Thr	Leu	Ser	Leu	Gln	Phe	Val	Asp	Ser	Ala	Ala	Asp	Met	Pro	Leu
			645						650			655			
Ala	Lys	Met	His	Gly	Ala	Phe	Ser	Thr	Asn	Val	Val	Ala	Ser	Lys	Glu
			660						665			670			
Leu	Gln	Gln	Pro	Gly	Ser	Ala	Arg	Ser	Thr	Arg	His	Leu	Glu	Ile	Glu
675						680						685			
Leu	Pro	Lys	Glu	Ala	Ser	Tyr	Gln	Glu	Gly	Asp	His	Leu	Gly	Val	Ile
690						695						700			
Pro	Arg	Asn	Tyr	Glu	Gly	Ile	Val	Asn	Arg	Val	Thr	Ala	Arg	Phe	Gly
705						710						715			
Leu	Asp	Ala	Ser	Gln	Gln	Ile	Arg	Leu	Glu	Ala	Glu	Glu	Glu	Lys	Leu
			725						730			735			
Ala	His	Leu	Pro	Leu	Ala	Lys	Thr	Val	Ser	Val	Glu	Glu	Leu	Leu	Gln
			740						745			750			
Tyr	Val	Glu	Leu	Gln	Asp	Pro	Val	Thr	Arg	Thr	Gln	Leu	Arg	Ala	Met
755						760						765			
Ala	Ala	Lys	Thr	Val	Cys	Pro	Pro	His	Lys	Val	Glu	Leu	Glu	Ala	Leu
770						775						780			

0050/50915

39

Leu	Glu	Lys	Gln	Ala	Tyr	Lys	Glu	Gln	Val	Leu	Ala	Lys	Arg	Leu	Thr	785	790	795	800
Met	Leu	Glu	Leu	Leu	Glu	Lys	Tyr	Pro	Ala	Cys	Glu	Met	Lys	Phe	Ser	805	810	815	
Glu	Phe	Ile	Ala	Leu	Leu	Pro	Ser	Ile	Arg	Pro	Arg	Tyr	Tyr	Ser	Ile	820	825	830	
Ser	Ser	Ser	Pro	Arg	Val	Asp	Glu	Lys	Gln	Ala	Ser	Ile	Thr	Val	Ser	835	840	845	
Val	Val	Ser	Gly	Glu	Ala	Trp	Ser	Gly	Tyr	Gly	Glu	Tyr	Lys	Gly	Ile	850	855	860	
Ala	Ser	Asn	Tyr	Leu	Ala	Glu	Leu	Gln	Glu	Gly	Asp	Thr	Ile	Thr	Cys	865	870	875	880
Phe	Ile	Ser	Thr	Pro	Gln	Ser	Glu	Phe	Thr	Leu	Pro	Lys	Asp	Pro	Glu	885	890	895	
Thr	Pro	Leu	Ile	Met	Val	Gly	Pro	Gly	Thr	Gly	Val	Ala	Pro	Phe	Arg	900	905	910	
Gly	Phe	Val	Gln	Ala	Arg	Lys	Gln	Leu	Lys	Glu	Gln	Gly	Gln	Ser	Leu	915	920	925	
Gly	Glu	Ala	His	Leu	Tyr	Phe	Gly	Cys	Arg	Ser	Pro	His	Glu	Asp	Tyr	930	935	940	
Leu	Tyr	Gln	Glu	Glu	Leu	Glu	Asn	Ala	Gln	Ser	Glu	Gly	Ile	Ile	Thr	945	950	955	960
Leu	His	Thr	Ala	Phe	Ser	Arg	Met	Pro	Asn	Gln	Pro	Lys	Thr	Tyr	Val	965	970	975	
Gln	His	Val	Met	Glu	Gln	Asp	Gly	Lys	Lys	Leu	Ile	Glu	Leu	Leu	Asp	980	985	990	
Gln	Gly	Ala	His	Phe	Tyr	Ile	Cys	Gly	Asp	Gly	Ser	Gln	Met	Ala	Pro	995	1000	1005	
Ala	Val	Glu	Ala	Thr	Leu	Met	Lys	Ser	Tyr	Ala	Asp	Val	His	Gln	Val	1010	1015	1020	
Ser	Glu	Ala	Asp	Ala	Arg	Leu	Trp	Leu	Gln	Gln	Leu	Glu	Glu	Lys	Gly	1025	1030	1035	1040
Arg	Tyr	Ala	Lys	Asp	Val	Trp	Ala	Gly								1045			